

VAN DER WAERDEN SPACES

MENACHEM KOJMAN

ABSTRACT. A topological space X is *van der Waerden* if for every sequence $(x_n)_n$ in X there exists a converging subsequence $(x_{n_k})_k$ so that $\{n_k : k \in \mathbb{N}\}$ contains arbitrarily long finite arithmetic progressions. Not every sequentially compact space is van der Waerden. The product of two van der Waerden spaces is van der Waerden.

The following condition on a Hausdorff space X is sufficient for X to be van der Waerden:

(*) The closure of every countable set in X is compact and first-countable.

A Hausdorff space X that satisfies (*) satisfies, in fact, a stronger property: for every sequence (x_n) in X :

(*) there exists $A \subseteq \mathbb{N}$ so that $(x_n)_{n \in A}$ is converging, and A contains arbitrarily long finite arithmetic progressions and sets of the form $FS(D)$ for arbitrarily large finite sets D .

There are nonmetrizable and non-compact spaces which satisfy (*). In particular, every sequence of ordinal numbers and every bounded sequence of real monotone functions on $[0, 1]$ satisfy (*).

1. INTRODUCTION

Let us recall Van der Waerden's Theorem. Let $\mathbb{N} = \{0, 1, \dots\}$ denote the set of natural numbers.

Definition 1. *A set A of natural numbers is called an AP set if it contains arbitrarily long finite arithmetic progressions.*

Theorem 2 (van der Waerden [9]). *If \mathbb{N} is partitioned to finitely many parts, then one of the parts is an AP set.*

An equivalent formulation of van der Waerden's theorem is that every sequence in a finite (Hausdorff) topological space has a converging subsequence whose set of indices contains arbitrarily long finite arithmetic progressions. In this paper I address the problem of which topological spaces satisfy this topological formulation of Van der Waerden's Theorems. The following condition (*) on a space X will be a sufficient condition in two different generalized van der Waerden type theorems (10 and 11 below):

1991 *Mathematics Subject Classification.* Primary: 05C55, 54F65; Secondary 03E05, 11P99, 26A48.

Key words and phrases. van der Waerden's Theorem, converging sequence, compactification, finite sums.

The author thanks Uri Abraham for many discussions that contributed to the development of this paper and thanks the referee for some constructive and helpful comments.

(*) The closure of every countable set in X is compact and first-countable.

2. VAN DER WAERDEN SPACES

Definition 3. 1. A topological space X is van der Waerden if for every sequence $(x_n)_n$ in X there exists a converging subsequence $(x_{n_k})_k$ so that $(n_k)_k$ is AP.
 2. A topological space X is strongly van der Waerden if for every AP set $A \subseteq \mathbb{N}$ and a sequence $(x_n)_{n \in A}$ in X there exists a converging subsequence $(x_n)_{n \in B}$ where $B \subseteq A$ is AP

Proposition 4. A space X is van der Waerden if and only if it is strongly van der Waerden.

Proof. Suppose $A \subseteq \mathbb{N}$ is AP and $f : A \rightarrow X$ is given. By thinning A out we may assume that $A = \bigcup C_n$ where each block C_n is an arithmetic progression, $\sup C_n < \min C_{n+1}$ and $|C_{n+1}| > (\max C_n)^2$.

Let $h : \mathbb{N} \rightarrow A$ be the increasing enumeration of A . Let $g : \mathbb{N} \rightarrow X$ be defined by $g(n) = f(h(n))$ for all $n \in \mathbb{N}$. Since X is van der Waerden, there exists an AP set $B \subseteq \mathbb{N}$ so that $g \upharpoonright B = (f \circ h) \upharpoonright B$ is converging. We claim that $h[B]$ is also AP, hence $f \upharpoonright h[B]$ is as required.

To see that $h[B]$ is AP, let k be arbitrary. Let $C = \{a_0, a_0 + d, \dots, a_0 + (k-1)d\} \subseteq B$ be an arithmetic progression of length k so that $k < a_0$. Observe that for each block C_n of A , $h[C] \cap C_n$ is an arithmetic progression.

Let C_{n_0} be the block in A to which $h(a_0 + d)$ belongs. Clearly k, a_0 and d are smaller than $\max C_{n_0}$, hence $a_0 + (k-1)d \leq \max C_{n_0} + (\max C_{n_0} - 1)(\max C_{n_0}) = (\max C_{n_0})^2 < |C_{n_0+1}|$. Since h is the increasing enumeration of $\bigcup C_n$, it holds that $h(a_0 + (k-1)d) \in C_{n_0} \cup C_{n_0+1}$, and consequently $h[C] \subseteq C_{n_0} \cup C_{n_0+1}$. Thus either $h[C] \cap C_{n_0}$ or $h[C] \cap C_{n_0+1}$ is an arithmetic progression of length $\geq (k-1)/2$ \square

Corollary 5. The product of two van der Waerden spaces is van der Waerden.

Proof. Suppose $Z = X \times Y$ where X, Y are van der Waerden and suppose $f : \mathbb{N} \rightarrow Z$ is given. Put $f(n) = (x_n, y_n)$. Since X is van der Waerden, there is an AP set $A \subseteq \mathbb{N}$ so that $(x_n)_{n \in A}$ is converging. Since Y is strongly van der Waerden, there is an AP set $B \subseteq A$ so that $(y_n)_{n \in B}$ is converging. Clearly, $(f(n))_{n \in B}$ is converging in Z . \square

Every van der Waerden space is surely sequentially compact. The next example shows that being van der Waerden is strictly stronger than being sequentially compact.

Theorem 6. There exists a Hausdorff, compact, sequentially compact, separable space which is first-countable at all points but one, which is not van der Waerden.

Proof. Let \mathcal{F} be a maximal member of $\{A : A \text{ is an infinite set of almost disjoint infinite subsets of } \mathbb{N} \text{ and each } A \in \mathcal{A} \text{ is not AP}\}$. Then in fact, \mathcal{F} is a maximal almost disjoint family. (If \mathcal{F} is not maximal as an almost disjoint family, fix some infinite $B \subseteq \mathbb{N}$ which has finite intersection with every $A \in \mathcal{F}$. By thinning out, B can be made non-AP.)

For every $A \in \mathcal{F}$, let $p_A \notin \mathbb{N}$ be a distinct point. Let $Y = \mathbb{N} \cup \{p_A : A \in \mathcal{F}\}$ and let $\mathcal{T} = \{B \subseteq Y : \text{for all } A \in \mathcal{F}, \text{ if } p_A \in B, \text{ then } A \setminus B \text{ is finite}\}$. Then with the

topology \mathcal{T} , Y is a locally compact Hausdorff space with \mathbb{N} as a discrete subspace. Let $X = Y \cup \{p\}$ be the one point compactification of Y .

X is compact, separable, and at every point except p , X is first countable. To check that X is sequentially compact, suppose that (x_n) is a sequence in X . If $\{x_n : n \in \mathbb{N}\}$ is finite, then $(x_n)_n$ has a constant subsequence. Else, we may assume, without loss of generality, that $(x_n)_n$ is 1-1 and that either $\{x_n : n \in \mathbb{N}\} \subseteq \mathbb{N}$ or $\{x_n : n \in \mathbb{N}\} \subseteq \{p_A : A \in \mathcal{F}\}$. In the first case, maximality of \mathcal{F} implies the existence of some $A \in \mathcal{F}$ and some subsequence $(x_{n_k})_k$ so that $x_{n_k} \in A$ for all k ; this subsequence converges to p_A . In the second case, $(x_n)_n$ itself converges to p . Thus X is sequentially compact.

To see that X is not van der Waerden, let $x_n = n$. Every convergent subsequence $(x_{n_k})_k$ satisfies that for some $A \in \mathcal{F}$, $\{n_k : k \in \mathbb{N}\} \setminus A$ is finite. By the choice of \mathcal{F} , $\{n_k : k \in \mathbb{N}\}$ is not AP. \square

The following is the well-known, equivalent, finite version of van der Waerden's theorem:

Theorem 7. *For all k and ℓ there exists n such that for every partition of $n = \{0, 1, \dots, n-1\}$ to ℓ parts, one of the parts contains an arithmetic progression of length k .*

One immediately sees that the finite version implies the following equivalent formulation of van der Waerden's theorem:

Theorem 8. *For every partition of an AP set to finitely many parts, one of the parts is AP.*

This formulation also follows directly from Proposition 4 applied to finite spaces.

Corollary 9. *The collection of all complements of non-AP sets is a uniform filter \mathcal{A} over \mathbb{N} which consists of AP sets. An ultrafilter U satisfies that every member of U is AP if and only if $\mathcal{A} \subseteq U$.*

Theorem 10. *Suppose a Hausdorff space X satisfies the following condition*

(*) *The closure of every countable set in X is compact and first-countable.*

Then X is van der Waerden.

Proof. Suppose that $(x_n)_{n \in \mathbb{N}} \subseteq X$ is given. Let $D = \text{cl}_X \{x_n : n \in \mathbb{N}\}$. Let U be an ultrafilter over \mathbb{N} which extends \mathcal{A} . Thus every set $A \in U$ is AP. Define an ultrafilter U' over D by $A \in U' \iff \{n : x_n \in A\} \in U$. By compactness (and Hausdorffness) of D , there is a unique point $x \in D$ with the property that every neighborhood (in D) of x belongs to U' . Fix, by first-countability, a decreasing neighborhood base $(u_k)_k$ at x . For every k , since $\{n : x_n \in u_k\} \in U$, the set $\{n : x_n \in u_k\}$ contains arbitrarily long arithmetic progressions. For each k , choose an arithmetic progression $C_k \subseteq \mathbb{N}$ of length k so that $\{x_n : n \in C_k\} \subseteq u_k$. The set $B = \bigcup_k C_k$ is AP and $(x_n)_{n \in B}$ converges to x . \square

2.1. A Strengthening. For a nonempty set $D \subseteq \mathbb{N}$ denote by $FS(D)$ the set of all *finite sums* over D . A finite sum is an integer of the form $\sum_{n \in \alpha} n$ where $\alpha \subseteq D$ is finite and nonempty. Call a subset $A \subseteq \mathbb{N}$ an IP set if there exists an infinite set $D \subseteq \mathbb{N}$ so that $FS(D) \subseteq A$.

Theorem 10 can be considerably strengthened using the technology of idempotents in $(\beta\mathbb{N}, +)$. $\beta\mathbb{N}$ is the set of all ultrafilters over \mathbb{N} and the operation of addition in \mathbb{N} can be extended to $\beta\mathbb{N}$ via the formula

$$(1) \quad A \in U + V \iff \{n : A - n \in U\} \in V$$

For a good exposition of this material see [2] (with the caution that they reverse the operation from the way it is described in (1)). An ultrafilter $U \in \beta\mathbb{N}$ is an *idempotent* if U is nonprincipal and $U + U = U$, and an idempotent U is a *minimal idempotent* if it belongs to a minimal right ideal in $(\beta\mathbb{N}, +)$. If U is an idempotent then every set $A \in U$ is IP and if U is a minimal idempotent then every set in U is AP (Theorems 4.40 and 14.1 in [2]).

Theorem 11. *Suppose a Hausdorff space X satisfies condition (*). Then for every sequence $(x_n)_{n \in \mathbb{N}} \subseteq X$ there exists a set $A \subseteq \mathbb{N}$ so that:*

1. $(x_n)_{n \in A}$ is converging;
2. A contains arbitrarily long arithmetic progressions
3. A contains $FS(D)$ for arbitrarily large finite sets D .

Proof. Let U be a minimal idempotent in $(\beta\mathbb{N}, +)$. Suppose $(x_n)_{n \in \mathbb{N}}$ is given, and let U' be the corresponding ultrafilter on $\text{cl}_X\{x_n : n \in \mathbb{N}\}$. Let $x \in \text{cl}_X\{x_n : n \in \mathbb{N}\}$ satisfy that every neighborhood of x belongs to U' , and let u_n be a decreasing neighborhood base at x . Since every set in U contains both arbitrarily long arithmetic progressions and an IP set, one can fix $FS(D_n)$ and an arithmetic progression C_n inside $\{k : x_k \in u_n\}$ with $|C_n|, |D_n| \geq n$. Now $\bigcup_n C_n \cup FS(D_n)$ is as required. \square

3. CONCLUDING REMARKS

The class of spaces which satisfy (*) includes all compact metric spaces, all compact linearly ordered topological spaces and every limit ordinal of uncountable cofinality. In particular,

Corollary 12. *Every sequence (α_n) of ordinal numbers contains a converging subsequence (α_{n_k}) so that $\{n_k : k \in \mathbb{N}\}$ contains arbitrarily long arithmetic progressions and $FS(D)$ for arbitrarily large finite sets D .*

The space ω_2 with the order topology is an example of a space that satisfies (*) which is neither first-countable nor compact (but all of its separable subspaces are metrizable). Helly's space of all monotone functions from $[0, 1]$ to $[0, 1]$ with the topology induced by pointwise convergence (see [3]) is an example of a separable, compact and first-countable space — which, therefore, satisfies (*) — which is not metrizable. The instance of Theorem 11 for this particular space gives:

Corollary 13. *Suppose that $f_n : [0, 1] \rightarrow [0, 1]$ is monotone for each n . Then there is pointwise converging subsequence (f_{n_k}) so that $\{n_k : k \in \mathbb{N}\}$ is AP and contains $FS(D)$ for arbitrarily large finite sets D .*

REFERENCES

- [1] Ronald L. Graham, Bruce L. Rothschild and Joel H. Spencer. Ramsey Theory, *John Wiley & Sons*, 1990.
- [2] N. Hindman and D. Strauss. Algebra in the Stone-Ćech compactification, *de Gruyter* Berlin, 1998.
- [3] Steen, Lynn Arthur and Seebach, Jr., J. Arthur. Counterexamples in topology, *Springer-Verlag*, New York, 1978

- [4] V. Bergelson. Ergodic Ramsey Theory — an update. *Ergodic theory and Z^d -actions*. London Math. Soc. Lecture Note series 228, 1–61, 1996
- [5] V. Bergelson and N. Hindman. Nonmetrizable topological dynamics and Ramsey Theory. *Trans. AMS*, 320:293–320, 1990.
- [6] N. Hindman. Finite sums from sequences within cells of partitions of \mathbb{N} . *J. Combinatorial Theory A*, 17:119–184, 1974.
- [7] N. Hindman. Ultrafilters and combinatorial number theory. *Number Theory Carbondale*, M. Nathanson, ed. *Lecture Notes in Math.*, 751, 1979.
- [8] N. Hindman. Partitions and sums and products of integers. *Trans. AMS*, 247, 227–245, 1979.
- [9] B. L. van der Waerden. Beweis eine Baudetschen Vermutung *Nieuw Arch. Wisk.*, 15:212–216, 1927.

DEPARTMENT OF MATHEMATICS, BEN GURION UNIVERSITY OF THE NEGEV, BEER SHEVA 84105,
ISRAEL

E-mail address: `kojman@math.bgu.ac.il`