

Colorful Strips

Greg Aloupis* · Jean Cardinal* · Sébastien Collette* · Shinji Imahori** · Matias Korman* · Stefan Langerman* · Oded Schwartz*** · Shakhar Smorodinsky† · Perouz Taslakian*

Received: date / Accepted: date

Abstract We study the following geometric hypergraph coloring problem: given a planar point set and an integer k , we wish to color the points with k colors so that any axis-aligned strip containing sufficiently many points contains all colors. We show that if the strip contains at least $2k-1$ points, such a coloring can always be found. In dimension d , we show that the same holds provided the strip contains at least $k(4\ln k + \ln d)$ points. We also consider the dual problem of coloring a given set of axis-aligned strips so that any sufficiently covered point in the plane is covered by k colors. We show that in dimension d the required coverage is at most $d(k-1) + 1$. This complements recent impossibility results on decomposition of strip coverings with arbitrary orientations.

From the computational point of view, we show that deciding whether a three-dimensional point set can be 2-colored so that any strip containing at least three points contains both colors is NP-complete. This shows a big contrast with the planar case, for which this decision problem is easy.

Keywords Hypergraph coloring · Covering decomposition · Lovász local lemma · Computational geometry

1 Introduction

There is a currently renewed interest in coloring problems on *geometric* hypergraphs, that is, set systems defined by geometric objects. This interest is motivated by applications to wireless and sensor networks [6]; conflict-free colorings [8], chromatic numbers [19], covering decompositions [17, 3], or polychromatic (*colorful*) colorings of geometric hypergraphs [4] have been extensively studied in this context.

Chargé de Recherches du FRS-FNRS.

Maître de Recherches du FRS-FNRS.

Université Libre de Bruxelles, CP212, Bld. du Triomphe, 1050 Brussels, Belgium. E-mail: {galoupis,jcardin,secollet,slanger,ptaslaki}@ulb.ac.be Supported by the Communauté française de Belgique - ARC. · Graduate School of Engineering, Nagoya University, Nagoya 464-8603, Japan E-mail: imahori@na.cse.nagoya-u.ac.jp · Departments of Mathematics, Technische Universität Berlin, 10623 Berlin, Germany. E-mail: odedsc@math.tu-berlin.de · Ben-Gurion University, Be'er Sheva 84105, Israel. E-mail: shakhar@math.bgu.ac.il

In this paper, we are interested in k -coloring finite point sets in \mathbb{R}^d so that any region bounded by two parallel axis-aligned hyperplanes, that contains at least some fixed number of points, also contains a point of each color.

An *axis-aligned strip*¹ is the area enclosed between two parallel axis-aligned hyperplanes. A k -coloring of a finite set assigns one of k colors to each element in the set. Let S be a k -colored set of points in \mathbb{R}^d . A strip is said to be *polychromatic* with respect to S if it contains at least one element of each color class. We define the function $p(k, d)$ as the minimum number for which there always exists a k -coloring of any point set in \mathbb{R}^d such that every strip containing at least $p(k, d)$ points is polychromatic. This is a particular case of the general framework proposed by Aloupis, Cardinal, Collette, Langerman, and Smorodinsky in [4].

Note that the problem does not depend on whether the strips are open or closed, since the problem can be seen in a purely combinatorial fashion: an axis-aligned strip isolates a subsequence of the points in sorted order with respect to one of the axes. Therefore, the only thing that matters is the order in which the points appear along each axis. We can thus rephrase our problem, considering d -dimensional points sets, as finding the minimum value $p(k, d)$ such that the following holds: For d permutations of a set of items S , it is always possible to color the items with k colors, so that in all d permutations every sequence of at least $p(k, d)$ contiguous items contains one item of each color.

We also study *circular* permutations, in which the first and the last elements are contiguous. We consider the problem of finding a minimum value $p'(k, d)$ such that, for any d circular permutations of a set of items S , it is possible to k -color the items so that in every permutation, every sequence of $p'(k, d)$ contiguous items contains all colors.

A restricted geometric version of this problem in \mathbb{R}^2 consists of coloring a point set S with respect to wedges. For our purposes, a wedge is any area delimited by two half-lines with common endpoint at one of d given apices. Each apex induces a circular ordering of the points in S . This is illustrated in Figure 1. We aim at coloring S so that any wedge containing at least $p'(k, d)$ points is polychromatic. In \mathbb{R}^2 , the non-circular case corresponds to wedges with apices at infinity, hence the circular case can be seen as a generalization.

We then study a dual version of the problem, in which a set of axis-aligned strips is to be colored so that sufficiently covered points are contained in strips from all color classes. For instance, in the planar case we study the following function $\overline{p}(k, d)$. Let H be a k -colored set of strips in \mathbb{R}^d . A point is said to be polychromatic with respect to H if it is contained in strips of all k color classes. The function $\overline{p}(k, d)$ is the minimum number for which there always exists a k -coloring of any set of strips in \mathbb{R}^d such that every point of \mathbb{R}^d contained in at least $\overline{p}(k, d)$ strips is polychromatic.

Note that the functions $p(k, d)$, $p'(k, d)$ and $\overline{p}(k, d)$ are monotone and non-decreasing (in particular, they all go to infinity when either k or d goes to infinity). Since we are interested in arbitrarily large pointsets, we always consider the set that we color to be “large enough” (that is, unbounded in terms of k).

Previous results. A *hypergraph* (S, R) is defined by a set S (called the *ground set*) and a set R of subsets of S . The main problem studied here is the coloring of geometric hypergraphs where the ground set S is a finite set of points, and the set of ranges R consists of all subsets of S that can be isolated by a single strip. In the dual case the ground set S is a finite set of geometric shapes and the ranges are points contained in the common intersection of a subset

¹ From here on, unless otherwise specified, a *strip* is always assumed to be axis-aligned




Fig. 1 Illustration of the definitions of $p(k,2)$ and $p'(k,2)$. On the left, points are 2-colored so that any axis-aligned strip containing at least three points is bichromatic. On the right, two points A and B define two circular permutations of the point set. In this case, we wish to color the points so that there is no long monochromatic subsequence in either of the two circular orderings.

of S . In some places in the literature, finite geometric hypergraphs are also referred to as *geometric range spaces*.

Several similar problems have been studied in this context [13,21,4], where the range space is not defined by strips, but rather by halfplanes, triangles, disks, pseudo-disks or translates of a centrally symmetric convex polygon. The problem was originally stated in terms of decomposition of c -covers (or f -fold coverings) in the plane: A c -cover of the plane by a convex body Q ensures that every point in the plane is covered by at least c translated copies of Q . In 1980, Pach [13] asked if, given Q , there exists a function $f(Q)$ such that every $f(Q)$ -cover of the plane can be decomposed into 2 disjoint 1-covers. A natural extension is to ask if given Q , there exists a function $f(k, Q)$ such that every $f(k, Q)$ -cover of the plane can be decomposed into k disjoint 1-covers. This corresponds to a k -coloring of the $f(k, Q)$ -cover, such that every point of the plane is polychromatic.

Partial answers to this problem are known: Pach [14] referenced an unpublished manuscript by Mani and Pach [11] showing that any 33-cover of the plane by unit disks can be decomposed into two 1-covers. This could imply that the function f exists for unit disks, but could still be exponential in k . Recently, Tardos and Tóth [21] proved that any 43-cover by translated copies of a triangle can be decomposed into two 1-covers. For the case of centrally symmetric convex polygons, Pach [15] proved that f is at most exponential in k . More than 20 years later, Pach and Tóth [17] improved this by showing that $f(k, Q) = O(k^2)$, and was afterwards Aloupis et al. [3] proved that $f(k, Q) = O(k)$. Recently, Gibson and Varadarajan [9] showed that the same property also holds for any arbitrary convex polygon Q .

On the other hand, for the range space induced by arbitrary disks, Mani and Pach [11] (see also [16]) proved that $f(2, Q)$ is unbounded: for any constant c , there exists a set of points that cannot be 2-colored so that all open disks containing at least c points are polychromatic. Pach, Tardos and Tóth [16] obtained a similar result for the range spaces induced by the family of either non-axis-aligned strips, or axis-aligned rectangles. Specifically, for any integer c there exist c -fold coverings with non-aligned strips that cannot be decomposed into two coverings (i.e., cannot be 2-colored). The previous impossibilities constitute our main

Table 1 Bounds on p , p' and \bar{p}

	$p(k, d)$	$p'(k, d)$	$\bar{p}(k, d)$
upper bound	$k(4 \ln k + \ln d)$ ($2k-1$ for $d=2$)	$k(4 \ln k + \ln d)$ ($2k$ for $d=2$)	$d(k-1) + 1$
lower bound	$2 \cdot \lceil \frac{(2d-1)k}{2d} \rceil - 1$		$\lfloor k/2 \rfloor d + 1$

motivation for introducing the problem of k -coloring axis-aligned strips, and strips with a bounded number of orientations.

Paper Organization. In Section 2 we give constructive upper bounds on the functions p and p' for $d = 2$. In Section 3 we consider higher-dimensional cases, as well as the computational complexity of finding a valid coloring. Section 4 concerns the dual problem of coloring strips with respect to points. Our lower and upper bounds are summarized in Table 1.

2 Axis-aligned strips and circular permutations for $d = 2$

We first consider upper bounds for the functions $p(k, 2)$ and $p'(k, 2)$.

2.1 Axis-aligned strips: Upper bound on $p(k, 2)$

We refer to a strip containing at least i points as an i -strip. Our goal is to show that for any integer k there is a constant $p(k, 2)$ such that any finite planar point set can be k -colored so that all $p(k, 2)$ -strips are polychromatic.

For $d = 2$, there is a reduction to the recently studied problem of 2-coloring graphs so that monochromatic components are small. Haxell et al. [10] proved that the vertices of any graph with maximum degree 4 can be 2-colored so that every monochromatic connected component has size at most 6. For a given finite point set S in the plane, let E be the set of all pairs of points $u, v \in S$ such that there is a strip containing only u and v . The graph $G = (S, E)$ has maximum degree 4, as it is the union of two paths. By the results of [10], G can be 2-colored so that every monochromatic connected component has size at most 6. In particular every path of size at least 7 contains points from both color classes. To finish the reduction argument one may observe that every strip containing at least 7 points corresponds to a path (of size at least 7) in G . We improve and generalize this first bound in the following.

Theorem 1 *For any finite planar set S and any integer k , S can be k -colored so that any $(2k-1)$ -strip is polychromatic. That is,*

$$p(k, 2) \leq 2k - 1.$$

Proof Let s_1, \dots, s_n be the points of S sorted by (increasing) x -coordinates and let $s_{\pi_1}, \dots, s_{\pi_n}$ be the sorting by y -coordinates. We first assume that k divides n , and later show how to remove the need for this assumption. Let V_x be the set of n/k disjoint contiguous k -tuples in s_1, \dots, s_n . Namely, $V_x = \{\{s_1, \dots, s_k\}, \{s_{k+1}, \dots, s_{2k}\}, \dots, \{s_{n-k+1}, \dots, s_n\}\}$. Similarly, let V_y be the k -tuples defined by $s_{\pi_1}, \dots, s_{\pi_n}$.

We define a bipartite multigraph $G = (V_x, V_y, E)$ as follows: For every pair of k -tuples $A \in V_x, B \in V_y$, we include an edge $e_s = \{A, B\} \in E$ if there exists a point s in both A and B . Note that an edge $\{A, B\}$ has multiplicity $|A \cap B|$ and that the number of edges $|E|$ is n . The multigraph G is k -regular because every k -tuple A contains exactly k points and every point $s \in A$ determines exactly one incident edge labeled e_s . It is well known that the chromatic index of any bipartite k -regular multigraph is k (and can be efficiently computed, see e.g., [1, 7]). Namely, the edges of such a multigraph can be partitioned into k perfect matchings. Let E_1, \dots, E_k be such a partition and $S_i \subset S$ be the set of labels of the edges of E_i . The sets S_1, \dots, S_k form a partition (i.e., a coloring) of S . We assign color i to the points of S_i .

We claim that this coloring ensures that any $(2k-1)$ -strip is polychromatic. Let h be a $(2k-1)$ -strip and assume without loss of generality that h is parallel to the y -axis. Then h contains at least one k -tuple $A \in V_x$. By the properties of the above coloring, the edges incident to A in G are colored with k distinct colors. Thus, the points that correspond to the labels of these edges are colored with k distinct colors, and h is polychromatic.

To complete the proof, we must handle the case where k does not divide n . Let $i = n \pmod{k}$. Let $Q = \{q_1, \dots, q_{k-i}\}$ be an additional set of $k-i$ points, all located to the right and above the points of S . We repeat our previous construction on $S \cup Q$. Now, any $(2k-1)$ -strip which is, say, parallel to the y -axis will also contain a k -tuple $A \in V_x$ disjoint from Q . Thus our arguments follow as before. \square

The proof of Theorem 1 is constructive and leads directly to an $O(n \log n)$ -time algorithm to k -color n points in the plane so that every $(2k-1)$ -strip is polychromatic. The algorithm is simple: we sort S , construct $G = (V_x, V_y, E)$, and color the edges of G with k colors. The time analysis is as follows: sorting takes $O(n \log n)$ time. Constructing G takes $O(n + |E|)$ time. As G has $\frac{2n}{k}$ vertices and is k -regular, it has n edges; so this step takes $O(n)$ time. Finding the edge-coloring of G takes $O(n \log n)$ time [1]. The total running time is therefore $O(n \log n)$.

2.2 Circular permutations: Upper bound on $p'(k, 2)$

We now consider the value of $p'(k, d)$. Given d circular permutations of a set S , we color S so that every sufficiently long subsequence in any of the circular permutations is polychromatic. The previous proof for $p(k, d) \leq 2k-1$ (Theorem 1) does not hold when we consider circular permutations. However, a slight modification provides the same upper bound, up to a constant term.

Theorem 2 $p'(k, 2) \leq 2k$

Proof If k divides n , we separate each circular permutation into n/k sets of size k . We define a multigraph, where the vertices represent the sets of k items, and there is an edge between two vertices if two sets share the same item. Trivially, this graph is k -regular and bipartite, and can thus be edge-colored with k colors. Each edge in this graph corresponds to one item in the permutation, thus each set of k items contains points of all k colors.

If k does not divide n , let $a = \lfloor n/k \rfloor$, and $b = n \pmod{k}$. If a divides b , we separate each of the two circular permutations into $2a$ sets, of alternating sizes k and b/a . Otherwise, the even sets will also alternate between size $\lceil b/a \rceil$ and $\lfloor b/a \rfloor$, instead of b/a . We extend both permutations by adding dummy items to each set of size less than k , so that we finally have only sets of size k . Dummy items appear in the same order in both permutations. We can now define the multigraph just as before.

If we remove the dummy nodes, we deduce a coloring for our original set. As each color appears in every set of size k , the length of any subsequence between two items of the same color is at most $2(k-1) + \lceil b/a \rceil$. Therefore, $p'(k, 2) \leq 2(k-1) + \lceil b/a \rceil + 1$.

Finally, if $n \geq k(k-1)$, then $a \geq k-1$, and $b \leq a$, we know that $\lceil b/a \rceil \leq 1$, and thus $p'(k, 2) \leq 2k$. \square

3 Higher dimensional strips

In this section we study the same problem for strips in higher dimensions. We provide upper and lower bounds on $p(k, d)$. We then analyze the coloring problem from a computational viewpoint, and show that deciding whether a given instance $S \subset \mathbb{R}^d$ can be 2-colored such that every 3-strip is polychromatic is NP-complete.

3.1 Upper bound on strip size, $p(k, d)$

Theorem 3 *Any finite set of points $S \subset \mathbb{R}^d$ can be k -colored so that every axis-aligned strip containing $k(4 \ln k + \ln d)$ points is polychromatic, that is,*

$$p(k, d) \leq k(4 \ln k + \ln d).$$

Proof The proof uses the probabilistic method. Let $\{1, \dots, k\}$ denote the set of k colors. We randomly color every point in S independently so that a point s gets color i with probability $\frac{1}{k}$ for $i = 1, \dots, k$. For a t -strip h , let \mathcal{B}_h be the “bad” event where h is not polychromatic. It is easily seen that $\Pr[\mathcal{B}_h] \leq k(1 - \frac{1}{k})^t$. Moreover, \mathcal{B}_h depends on at most $(d-1)t^2 + 2t - 2$ other events. Indeed, \mathcal{B}_h depends only on t -strips that share points with h . Assume without loss of generality that h is orthogonal to the x_1 axis. Then \mathcal{B}_h has a non-empty intersection with at most $2(t-1)$ other t -strips which are orthogonal to the x_1 axis. For each of the other $d-1$ axes, h can intersect at most t^2 t -strips since every point in h can belong to at most t other t -strips.

By the Lovász Local Lemma, (see, e.g., [2]) we have that if t satisfies

$$e \cdot ((d-1) \cdot t^2 + 2t - 1) \cdot k \left(1 - \frac{1}{k}\right)^t < 1$$

(where e is the basis of the natural logarithm), then

$$\Pr \left[\bigwedge_{|h|=t} \overline{\mathcal{B}_h} \right] > 0.$$

In particular, this means that there exists a k -coloring for which every t -strip is polychromatic. It can be verified that $t = k(4 \ln k + \ln d)$ satisfies the condition. \square

The proof of Theorem 3 is non-constructive. However, we can use known algorithmic versions of the Local Lemma (see for instance [12]) to obtain a constructive proof. Also note that Theorem 3 holds in the more general case where the strips are not necessarily axis-aligned. In fact, one can have a total of d arbitrary strip orientations in some fixed arbitrary dimension and the proof will hold verbatim. Finally, we note that the same proof also works for the case of circular permutations, yielding the same upper bound:

Theorem 4 $p'(k, d) \leq k(4 \ln k + \ln d)$

3.2 Lower bound on $p(k, d)$

We first introduce a well-known result on the decomposition of complete graphs:

Lemma 1 *The edges of K_{2h} can be decomposed into h pairwise edge-disjoint Hamiltonian paths.*

This result follows from a special case of the Oberwolfach problem [5]. An explicit proof of this lemma can also be found in [20].

Note that if the vertices of K_{2h} are labeled $V = \{1, \dots, 2h\}$, each path can be seen as a permutation of $2h$ elements. Using Lemma 1 we obtain:

Theorem 5 *For any fixed dimension d and number of colors k , let $s = \left\lceil \frac{(2d-1)k}{2d} \right\rceil - 1$. Then,*

$$p'(k, d) \geq p(k, d) \geq 2s + 1.$$

Proof The first inequality comes from the fact that any polychromatic coloring with respect to circular permutations is also polychromatic with respect to strips. We now focus on showing the second inequality: let $\sigma_1, \dots, \sigma_d$ be any decomposition of K_{2d} into d paths: we construct the set $P = \{p_i | 0 \leq i \leq 2d\}$, where $p_i = (\sigma_1(i), \dots, \sigma_d(i))$. Note that the ordering of P , when projected to the i -th axis, gives permutation σ_i . Since the elements σ decompose K_{2d} , in particular for any $i, j \leq 2d$ there exists a permutation in which i and j are adjacent.

We replace each point p_i by a set A_i of s points arbitrarily close to p_i . By construction, for any $i, j \leq 2d$, there exists a $2s$ -strip containing exactly $A_i \cup A_j$. Consider any possible coloring of the sets A_i : since $|A_i| = s$ and we are using k colors, there are at least $k - s$ colors not present in any set A_i .

Since $\left\lceil \frac{(2d-1)k}{2d} \right\rceil - 1 < \frac{(2d-1)k}{2d}$, we conclude that $k - s > k - \frac{(2d-1)k}{2d} = k/2d$. That is, each set is missing strictly more than $k/2d$ colors. By the pigeonhole principle, there exist i and j such that the set $A_i \cup A_j$ is missing a color (otherwise there would be more than k colors). In particular, the strip that contains set $A_i \cup A_j$ is not polychromatic, thus the theorem is shown.

We gave a set of bounded size $n = 2d$ reaching the lower bound, but we can easily create larger sets reaching the same bound: we can add as many dummy points as needed at the end of every permutation, which does not decrease the value of $p(k, d)$. \square

Note that, asymptotically speaking, the lower bound does not depend on d . However, by the negative result of [16], we know that $p(k, d) \rightarrow \infty$ when $d \rightarrow \infty$.

3.3 Computational complexity

In Section 2, we provided an algorithm that finds a k -coloring such that every planar $(2k-1)$ -strip is polychromatic. Thus for $d=2$ and $k=2$, this yields a 2-coloring such that every 3-strip is polychromatic.

Note that in this case $p(2, 2) = 3$, but the minimum required size of a strip for a given instance can be either 2 or 3. Testing if it is equal to 2 is easy: we can simply alternate the colors in the first permutation, and check if they also alternate in the other. Hence the problem of minimizing the size of the largest monochromatic strip on a given instance is polynomial for $d = 2$ and $k = 2$. We now show that it becomes NP-hard for $d > 2$ and $k = 2$. The same problem for $k > 2$ is left open.

Theorem 6 *The following problem is NP-complete:*

Input: 3 permutations π_1, π_2, π_3 of an n -element set S .

Question: *Is there a 2-coloring of S , such that every 3 elements of S that are consecutive according to one of the permutations are not monochromatic?*

Proof We show a reduction from NAE 3SAT (not-all-equal 3SAT) which is the following NP-complete problem [18]:

Input: A 3-CNF Boolean formula Φ .

Question: Is there a NAE assignment to Φ ? An assignment is called NAE if every clause has at least one literal assigned True and at least one literal assigned False.

We first transform Φ into another instance Φ' in which all variables are non-negated (i.e., we make the instance monotone). We then show how to realize Φ' using three permutations π_1, π_2, π_3 .

To transform Φ into Φ' , for each variable x , we first replace the i th occurrence of x in its positive form by a variable x_i , and the i th occurrence of x in its negative form by x'_i . The index i varies between 1 and the number of occurrences of each form (the maximum of the two). We also add the following *consistency-clauses*, for each variable x and for all i :

$$(Z_i^x, x_i, x'_i), (x_i, x'_i, Z_{i+1}^x), (x_i, Z_i^{x'}, x'_i), (Z_i^x, Z_i^{x'}, Z_{i+1}^x) \\ (x'_i, Z_{i+1}^x, x_{i+1}), (Z_i^{x'}, x'_i, x_{i+1}), (x'_i, x_{i+1}, Z_{i+1}^{x'}), (Z_i^{x'}, Z_{i+1}^x, Z_{i+1}^{x'})$$

where Z_i^x and $Z_i^{x'}$ are new variables. This completes the construction of Φ' . Note that Φ' is monotone, as every negated variable has been replaced.

Moreover, Φ' has a NAE assignment if and only if Φ has a NAE assignment. To see this, note that a NAE assignment for Φ can be translated to a NAE assignment to Φ' as follows: for every variable x of Φ and every i , set $x_i \equiv x$, $x'_i \equiv \bar{x}$, $Z_i^x \equiv True$, $Z_i^{x'} \equiv False$.

On the other hand, if Φ' has a NAE assignment, then, by the consistency clauses, the variables in Φ' corresponding to any variable x of Φ are assigned a consistent value. Namely, for every i, j we have $x_i = x_j$ and $x_i \neq x'_i$. This assignment naturally translates to a NAE assignment for Φ , by setting $x \equiv x_1$.

We next show how to realize Φ' by a set S and three permutations π_1, π_2, π_3 . The elements of the set S are the variables of Φ' , together with some additional elements that are described below. Permutation π_1 realizes the clauses of Φ' corresponding to the original clauses of Φ , while π_2 and π_3 realize the consistency clauses of Φ' .

The additional elements in S are clause elements (two elements c_{2j-1} and c_{2j} for every clause j of Φ) and *dummy elements* \star (the dummy elements are not indexed for the ease of presentation, but they appear in the same order in all three permutations).

Permutation π_1 encodes the clauses of Φ' corresponding to original clauses of Φ as follows (note that all these clauses involve different variables). For each such clause (u, v, w) , permutation π_1 contains the following sequence:

$$c_{2j-1}, u, v, w, c_{2j}, \star, \star$$

At the end of π_1 , for every variable x of Φ' we have the sequence:

$$Z_1^x, Z_1^{x'}, Z_2^x, Z_2^{x'}, Z_3^x, Z_3^{x'}, \dots, \star, \star$$

Permutation π_2 contains, for every variable x of Φ , the sequences:

$$Z_1^x, x_1, x'_1, Z_2^x, x_2, x'_2, Z_3^x, x_3, x'_3, Z_4^x, \dots \quad \text{and} \quad \star, \star, Z_1^{x'}, \star, \star, Z_2^{x'}, \star, \star, Z_3^{x'}, \dots, \star, \star$$

At the end of π_2 we have the clause-elements and remaining dummy elements:

$$\star, \star, c_1, \star, \star, c_2, \star, \star, c_3, \dots$$

Similarly, permutation π_3 contains, for every variable x of Φ , the sequences:

$$x_1, Z_1^x, x'_1, x_2, Z_2^x, x'_2, x_3, Z_3^x, x'_3, \dots \quad \text{and} \quad \star, \star, Z_1^x, \star, \star, Z_2^x, \star, \star, Z_3^x, \dots \star, \star$$

and at the end of π_3 we have the clause-elements and remaining dummy elements:

$$\star, \star, c_1, \star, \star, c_2, \star, \star, c_3, \dots$$

This completes the construction of S and π_1, π_2, π_3 . Note that for every clause of Φ' (whether it is derived from Φ or is a consistency clause), the elements corresponding to its three variables appear in sequence in one of the three permutations. Therefore, if there is a 2-coloring of S , such that every 3 elements of S that are consecutive according to one of the permutations are not monochromatic, then there is a *NAE* assignment to Φ' : each variable of Φ' is assigned True if its corresponding element is colored '1', and False otherwise.

For the other direction, consider a *NAE* assignment for Φ' . Observe that there is always a solution where Z_i^x and $Z_i^{x'}$ are assigned opposite values. Then assign color '1' to elements corresponding to variables assigned with True, and assign color '0' to elements corresponding to variables assigned with False. For the clause elements c_{2j-1} and c_{2j} appearing in the subsequence $c_{2j-1}, u, v, w, c_{2j}$, assign to c_{2j-1} the color opposite to u , and to c_{2j} the color opposite to w . Finally, assign colors '0' and '1' to each pair of consecutive dummy elements, respectively. It can be verified that there is no monochromatic consecutive triple in any permutation. \square

Approximation. Note that the general minimization problem (find a k -coloring that minimizes the number of required points) can be approximated using the constructive version of the Lovász Local Lemma (as mentioned in Section 3.1). Since k is a trivial lower bound for any problem instance, this guarantees an approximation factor of $O(\log k + \log d)$. In particular, there exists a constant factor approximation for the problem introduced in Theorem 6 (since it fixes $d = 3$ and $k = 2$).

4 Coloring strips

In this section we prove that any finite set of strips in \mathbb{R}^d can be k -colored so that every "deep" point is polychromatic. For a given set of strips (or intervals, if $d = 1$), we say that a point is i -deep if it is contained in at least i of the strips. We begin with the following easy lemma:

Lemma 2 *Let \mathcal{I} be a finite set of intervals. Then for every k , \mathcal{I} can be k -colored so that every k -deep point is polychromatic, while any point covered by fewer than k intervals will be covered by distinct colors.*

Proof We use induction on $|\mathcal{I}|$. Let I be the interval with the leftmost right endpoint. By induction, the intervals in $\mathcal{I} \setminus \{I\}$ can be k -colored with the desired property. Sort the intervals intersecting I according to their left endpoints and let I_1, \dots, I_{k-1} be the first $k-1$ intervals in this order. It is easily seen that coloring I with a color distinct from the colors of those $k-1$ intervals produces a coloring with the desired property, and hence a valid coloring. \square

Theorem 7 For any d and k , one can k -color any set of axis-aligned strips in \mathbb{R}^d so that every $d(k-1)+1$ -deep point is polychromatic. That is,

$$\bar{p}(k, d) \leq d(k-1) + 1.$$

Proof We start by coloring the strips parallel to any given axis x_i ($i = 1, \dots, d$) separately using the coloring described in Lemma 2. We claim that this procedure produces a valid polychromatic coloring for all $d(k-1)+1$ -deep points. Indeed assume that a given point s is $d(k-1)+1$ -deep and let $H(s)$ be the set of strips covering s . Since there are d possible orientations for the strips in $H(s)$, by the pigeonhole principle at least k of the strips in $H(s)$ are parallel to the same axis. Then by property of the coloring of Lemma 2, $H(s)$ is polychromatic. \square

The above proof is constructive. By sorting the intervals that correspond to any of the given directions, one can easily find a coloring in $O(n \log n)$ time.

We now give a lower bound on $\bar{p}(k, d)$. For that, we define $2d$ strips as follows: strip s_{2i} is defined as $0 < x_i < 2$ (where x_i is the coordinate of i -th dimension). Analogously, we define strip s_{2i+1} as $1 < x_i < 3$. The main property of these strips is that we can always find a point covered by any subset of the $2d$ strips:

Lemma 3 For any $I \subseteq \{1, \dots, 2d\}$, there exists a point p_I such that $p_I \in s_i \Leftrightarrow i \in I, \forall i \leq 2d$.

Proof Note that whether or not point p_I is covered by strips s_{2i} or s_{2i+1} only depends in the i -th coordinate of p_I . Thus, we define the i -th coordinate of point p_I as follows:

- 1 if $2i \in I$ but $2i+1 \notin I$
- 2 if both $2i, 2i+1 \in I$
- 3 if both $2i, 2i+1 \notin I$

Since the choice is independent on each dimension, the construction of p_I is valid and is only covered by strips in I . \square

We use these strips to find a lower bound on $\bar{p}(k, d)$ as follows:

Theorem 8 For any fixed dimension d and integer k , it holds that

$$\bar{p}(k, d) > \lfloor k/2 \rfloor d + 1.$$

Proof Consider the $2d$ strips $\{s_i\}_{i \in [2d]}$ defined above. We replace each strip s_i with a cluster of $\lfloor k/2 \rfloor$ overlapping strips $\{s_{i,j}\}_{j \in [\lfloor k/2 \rfloor]}$, so that a point p_I defined in Lemma 3 is in strip $s_{i,j}$ if and only if it is in s_i . This can be obtained, say, by perturbing a boundary of the strip around $x_i = 0$ (or around $x_i = 3$).

Consider any coloring of the above problem instance with at most $\lfloor k/2 \rfloor d$ colors. As in the proof of Theorem 5, we can use the pigeonhole principle and the handshake lemma to conclude that there is a color that is missing in at least $\lceil k/2 \rceil 2d/k \geq d$ clusters. Let I be the set of at least d indices whose clusters are missing the same color. By Lemma 3, point p_I is covered only by the strips in clusters C_i , for all $i \in I$. In particular, p_I is at least $\lfloor k/2 \rfloor d$ -deep and not colorful. \square

Acknowledgements This research was initiated during the WAFOL'09 workshop at Université Libre de Bruxelles (U.L.B.), Brussels, Belgium. The authors want to thank all other participants, and in particular Erik D. Demaine, for helpful discussions. Also, the authors would like to thank the anonymous referees for pointing out useful references.

References

1. Alon, N.: A simple algorithm for edge-coloring bipartite multigraphs. *Inf. Proc. Lett.* **85**(6), 301–302 (2003). DOI [http://dx.doi.org/10.1016/S0020-0190\(02\)00446-5](http://dx.doi.org/10.1016/S0020-0190(02)00446-5)
2. Alon, N., Spencer, J.: *The Probabilistic Method*, 2nd edition. John Wiley (2000)
3. Aloupis, G., Cardinal, J., Collette, S., Langerman, S., Orden, D., Ramos, P.: Decomposition of multiple coverings into more parts. In: *Proceedings of the ACM-SIAM Symposium on Discrete Algorithms (SODA'09)* (2009)
4. Aloupis, G., Cardinal, J., Collette, S., Langerman, S., Smorodinsky, S.: Coloring geometric range spaces. *Discrete & Computational Geometry* **41**(2), 348–362 (2009)
5. B. Alspach: The wonderful walecki construction. *Bull. Inst. Combin. Appl.* **52**, 7–20 (2008)
6. Buchsbaum, A., Efrat, A., Jain, S., Venkatasubramanian, S., Yi, K.: Restricted strip covering and the sensor cover problem. In: *ACM-SIAM Symposium on Discrete Algorithms (SODA'07)* (2007)
7. Cole, R., Ost, K., Schirra, S.: Edge-coloring bipartite multigraphs in $O(E \log D)$ time. *Combinatorica* **21**(1), 5–12 (2001)
8. Even, G., Lotker, Z., Ron, D., Smorodinsky, S.: Conflict-free colorings of simple geometric regions with applications to frequency assignment in cellular networks. *SIAM Journal on Computing* **33**(1), 94–136 (2004)
9. Gibson, M., Varadarajan, K.R.: Decomposing coverings and the planar sensor cover problem. *CoRR abs/0905.1093* (2009). Informal publication (to appear in FOCS'09)
10. Haxell, P., Szabó, T., Tardos, G.: Bounded size components: partitions and transversals. *J. Comb. Theory Ser. B* **88**(2), 281–297 (2003). DOI [http://dx.doi.org/10.1016/S0095-8956\(03\)00031-5](http://dx.doi.org/10.1016/S0095-8956(03)00031-5)
11. Mani, P., Pach, J.: Decomposition problems for multiple coverings with unit balls (1986). Manuscript
12. Moser, R.A., Tardos, G.: A constructive proof of the general Lovász local lemma. *J. ACM* **57**(2) (2010)
13. Pach, J.: Decomposition of multiple packing and covering. In: *2. Kolloq. über Diskrete Geom.*, pp. 169–178. Inst. Math. Univ. Salzburg (1980)
14. Pach, J.: Decomposition of multiple packing and covering. In: *2. Kolloquium Über Diskrete Geometrie*, pp. 169–178. Inst. Math. Univ. Salzburg, Salzburg (1980)
15. Pach, J.: Covering the plane with convex polygons. *Discrete & Computational Geometry* **1**, 73–81 (1986)
16. Pach, J., Tardos, G., Tóth, G.: Indecomposable coverings. In: *The China–Japan Joint Conference on Discrete Geometry, Combinatorics and Graph Theory (CJCDGCGT 2005)*, Lecture Notes in Computer Science, pp. 135–148 (2007)
17. Pach, J., Tóth, G.: Decomposition of multiple coverings into many parts. In: *Proc. of the ACM Symposium on Computational Geometry*, pp. 133–137 (2007)
18. Schaefer, T.J.: The complexity of satisfiability problems. In: *STOC*, pp. 216–226. ACM (1978)
19. Smorodinsky, S.: On the chromatic number of some geometric hypergraphs. *SIAM Journal on Discrete Mathematics* **21**(3), 676–687 (2007)
20. Stanton, R.G., Cowan, D.D., James, L.O.: Some results on path numbers. In: *Louisiana Conference on Combin., Graph Theory and Computing* (1970)
21. Tardos, G., Tóth, G.: Multiple coverings of the plane with triangles. *Discrete & Computational Geometry* **38**(2), 443–450 (2007)