

Improved generalization bounds for robust learning

Idan Attias

Aryeh Kontorovich

Department of Computer Science

Ben-Gurion University of the Negev, Beer Sheva, Israel

IDANATTI@POST.BGU.AC.IL

KARYEH@CS.BGU.AC.IL

Yishay Mansour

Blavatnik School of Computer Science, Tel Aviv University, Tel Aviv Israel

and Google Research, Tel Aviv

MANSOUR.YISHAY@GMAIL.COM

Editor:

Abstract

We consider a model of robust learning in an adversarial environment. The learner gets uncorrupted training data with access to possible corruptions that may be effected by the adversary during testing. The learner's goal is to build a robust classifier that would be tested on future adversarial examples. We use a zero-sum game between the learner and the adversary as our game theoretic framework. The adversary is limited to k possible corruptions for each input. Our model is closely related to the adversarial examples model of [Schmidt et al. \(2018\)](#); [Madry et al. \(2017\)](#).

Our main results consist of generalization bounds for the binary and multi-class classification, as well as the real-valued case (regression). For the binary classification setting, we both tighten the generalization bound of [Feige, Mansour, and Schapire \(2015\)](#), and also are able to handle an infinite hypothesis class \mathcal{H} . The sample complexity is improved from $\mathcal{O}(\frac{1}{\epsilon^4} \log(\frac{|\mathcal{H}|}{\delta}))$ to $\mathcal{O}(\frac{1}{\epsilon^2} (k \log(k) VC(\mathcal{H}) + \log \frac{1}{\delta}))$. Additionally, we extend the algorithm and generalization bound from the binary to the multiclass and real-valued cases. Along the way, we obtain results on fat-shattering dimension and Rademacher complexity of k -fold maxima over function classes; these may be of independent interest.

For binary classification, the algorithm of [Feige et al. \(2015\)](#) uses a regret minimization algorithm and an ERM oracle as a blackbox; we adapt it for the multi-class and regression settings. The algorithm provides us with near optimal policies for the players on a given training sample.

Keywords: Robust Learning, Adversarial Learning, Generalization Bounds, Rademacher Complexity, Fat-Shattering Dimension, Zero-Sum Game

1. Introduction

We study the classification and regression problems in a setting of adversarial examples. This setting is different from standard supervised learning in that examples, both at training and testing time, may be corrupted in an adversarial manner to disrupt the learner's performance. This challenge to design reliable robust models gained significant attention as standard supervised learning methods have shown vulnerability, and is named *adversarial examples*. We study the adversarially robust learning paradigm from a generalization point of view and concentrate on the case of having adversarial examples at test time.

We consider the following robust learning framework for multi-class and real valued functions of [Feige et al. \(2015\)](#). There is an unknown distribution over the uncorrupted inputs domain. The learner receives a labeled uncorrupted sample (the labels can be categorical or real valued) and has access during the training phase to all possible corruptions that the adversary might effect. The learner selects a hypothesis from a fixed hypothesis class (in our case, weighted average of classifiers from base class \mathcal{H}) that gives a prediction for a corrupted input. The learner's accuracy is measured

by predicting the true label of the uncorrupted input while they observe only the corrupted input during test time. Thus, their goal is to find a policy that is immune to those corruptions. The adversary is capable of corrupting each future input, but there are only k possible corruptions for each input. This leads to a game theoretic framework of a zero-sum game between the learner and the adversary.

The model is closely related to the one suggested by [Schmidt et al. \(2018\)](#); [Madry et al. \(2017\)](#) and common robust optimization approaches ([Ben-Tal et al., 2009](#)), which deal with bounded worst-case perturbations (under l_∞ norm) on the samples. In this work we do not assume any metric for the corruptions, the adversary can map an input from the sample space to any other space, but is limited with finite possible corruptions for each input.

Our focus is on adversarial examples during testing time. The training data is clean, but we take into consideration all possible corruptions when we build the robust classifier. Thus, we extend the ERM paradigm by using adversarial training techniques instead of merely find a hypothesis that minimizes the empirical risk. Contrary to “standard” learning, ERM often does not yield models that are robust to adversarially corrupted examples ([Goodfellow et al., 2015](#); [Kurakin et al., 2017](#); [Moosavi-Dezfooli et al., 2016](#); [Tramèr et al., 2017](#)). Another point of interest is when the training data is adversarially corrupted beforehand, without any option to access the uncorrupted sample.

Studying worst-case (adversarial) corruptions is interesting for a couple of reasons. First, it models situations where the corruption occurs maliciously (such as a spammer who tailors messages to avoid a spam detector) and not merely as a result of random noise. Additionally, a robust classifier that is successful against adversarial corruption extends to less adversarial settings.

Our main results are generalization bounds for all settings. For the binary classification setting, we improve the generalization bound given in [Feige et al. \(2015\)](#). We generalize to the case of weighted average of hypotheses from \mathcal{H} when \mathcal{H} is not necessarily finite. The sample complexity has been improved from $\mathcal{O}(\frac{1}{\epsilon^4} \log(\frac{|\mathcal{H}|}{\delta}))$ to $\mathcal{O}(\frac{1}{\epsilon^2} (k \log(k) \text{VC}(\mathcal{H}) + \log \frac{1}{\delta}))$. The core of all proofs is a bound on the Rademacher complexity of the k -fold maximum of the convex hull of \mathcal{H} (the learner’s classifier generated by the algorithm is a member of this set). The k -fold maximum captures the k possible corruptions for each input. In the regression case we provide a tight bound on the fat shattering dimension of k -fold maximum class and bound the fat shattering dimension of L_1 and L_2 loss classes.

For our algorithm, we employ a regret minimization algorithm proposed for binary classification by [Feige et al. \(2015\)](#) for computing near optimal policies for the players on the training data. We adapt it multiclass classification and regression as well. The algorithm is a variant of the algorithm found in [Cesa-Bianchi et al. \(2005\)](#) and based on the ideas of [Freund and Schapire \(1999\)](#). An ERM (empirical risk minimization) oracle is used multiple times to return a hypothesis that minimizes the error rate on a given sample, while weighting samples differently every time. The learner uses a fixed hypotheses class \mathcal{H} and their chosen hypothesis is a convex combination of hypotheses from \mathcal{H} .

1.1 Related work

The most related work studying robust learning with adversarial examples are [Schmidt et al. \(2018\)](#); [Madry et al. \(2017\)](#). Their model deals with bounded worst-case perturbations (under l_∞ norm) on the samples. This is slightly different from our model as we mentioned above.

All of our results based on a robust learning model for binary classification suggested by [Feige et al. \(2015\)](#). In this work, we provide a tighter generalization bound and expand on multiclass classification and regression settings. The works of [Mansour et al. \(2015\)](#); [Feige et al. \(2015, 2018\)](#)

consider *robust inference* for the binary and multi-class case. The robust inference model assumes that the learner knows both the distribution and the target function, and the main task is given a corrupted input, derive in a computationally efficient way a classification which will minimize the error. In this work we consider only the learning setting, where the learner has only access to an uncorrupted sample, and need to approximate the target function on possibly corrupted inputs, using a restricted hypothesis class \mathcal{H} .

The work of [Globerson and Roweis \(2006\)](#) and its extensions [Teo et al. \(2008\)](#); [Dekel and Shamir \(2008\)](#) discuss a robust learning model where an uncorrupted sample is drawn from an unknown distribution, and the goal is to learn a linear classifier that would be able to overcome missing attributes in future test examples. They discuss both the static model (where the set of missing attributes is selected independently from the uncorrupted input) and the dynamic model (where the set of missing attributes may depend on the uncorrupted input). The model we use ([Feige et al., 2015](#)) extends the robust learning model to handle corrupted inputs (and not only missing attributes) and an arbitrary hypothesis class (rather than only linear classifiers).

There is a vast literature in statistics, operation research and machine learning regarding various noise models. Typically, most noise models assume a random process that generates the noise. In computational learning theory, popular noise models include random classification noise ([Angluin and Laird, 1988](#)) and malicious noise ([Valiant, 1985](#); [Kearns and Li, 1993](#)). In the malicious noise model, the adversary gets to arbitrarily corrupt some small fraction of the examples; in contrast, in our model the adversary can always corrupt every example, but only in a limited way.

Other motivating applications such as spam messages detection, web spam detection, computer intrusion detection, fraud detection, network failure detection, noisy bio-sensors measurements, and many more can be found in [Laskov and Lippmann \(2010\)](#).

1.2 The structure of the paper

The structure of this paper is as follows: Section 2 discusses the model in detail. Section 3 contains relevant definitions and notations. Section 4 is the learning algorithm, and Sections 5, 6 and 7 contain the generalization bounds for binary and multiclass classification and regression.

2. Model

There is some unknown distribution D over a finite domain \mathcal{X} of uncorrupted examples and a finite domain of corrupted examples \mathcal{Z} , possibly the same as \mathcal{X} . We work in deterministic scenario, where there is some concept class C such that $c \in C$ has domain \mathcal{X} and range \mathcal{Y} that can be $\{1, \dots, l\}$ or \mathbb{R} . There is some unknown target function $c^* \in C$ maps an uncorrupted example to its label.

The adversary is able to corrupt an input by mapping an uncorrupted input $x \in \mathcal{X}$ to a corrupted one $z \in \mathcal{Z}$. There is a mapping ρ which for every $x \in \mathcal{X}$ defines a set $\rho(x) \subseteq \mathcal{Z}$, such that $|\rho(x)| \leq k$. The adversary can map an uncorrupted input x to any corrupted input $z \in \rho(x)$. We assume that the learner has an access to $\rho(\cdot)$ during the training phase.

There is a fixed hypothesis class \mathcal{H} of hypothesis $h : \mathcal{Z} \mapsto \mathcal{Y}$ over corrupted inputs. The learner observes an uncorrupted sample $S_u = \{\langle x_1, c^*(x_1) \rangle, \dots, \langle x_m, c^*(x_m) \rangle\}$, where x_i is drawn i.i.d. from D , and needs to select a hypothesis $h \in \mathcal{H}$ (or a mixture of hypotheses $h \in \text{conv}(\mathcal{H})$) from the hypothesis class \mathcal{H} . The main difference from the classical learning models is that the learner will be tested on adversarially corrupted inputs $z \in \rho(x)$ and when selecting an hypothesis $h \in \mathcal{H}$ this need to be taken into consideration.

The basic scenario is as follows. First, an uncorrupted input \mathcal{X} is selected using D . Then, the adversary, given \mathcal{X} , selects some $z \in \rho(x)$. The learner observes a corrupted input \mathcal{Z} and outputs a mixture of labels, as dictated by $h \in \text{conv}(\mathcal{H})$, for predicting the value of $c^*(x)$. Finally, the learner incurs a loss. We use the zero-one loss to measure the quality of the classification, i.e., $\text{loss}(h(z), y) = I_{\{h(z) \neq y\}}$. In the regression case we use l_1 and l_2 loss functions, i.e., $\text{loss}(h(z), y) = |h(z) - y|^p$, for $p = 1, 2$.

The goal of the learner is to minimize the expected loss, while the adversary would like to maximize it. This defines a zero-sum game which has a value v which is the learner's error rate. We say that the learner's hypothesis is ϵ -optimal if it guarantees a loss which is at most $v + \epsilon$, and the adversary policy is ϵ -optimal if it guarantees a loss which is at least $v - \epsilon$. We refer to a 0-optimal policy as an optimal policy.

Formally, the error (risk) of the learner when selecting a hypothesis $h \in \mathcal{H}$ is

$$R(h) = \mathbb{E}_{x \sim D} [\max_{z \in \rho(x)} \text{loss}(h(z), c^*(x))],$$

and his goal is to choose $h \in \mathcal{H}$ to minimize the loss

$$\min_{h \in \mathcal{H}} R(h) = \min_{h \in \mathcal{H}} \mathbb{E}_{x \sim D} [\max_{z \in \rho(x)} \text{loss}(h(z), c^*(x))].$$

The uncorrupted training data learning algorithm (UTD-learning algorithm) receives an uncorrupted sample S_u and outputs a hypothesis $h \in \mathcal{H}$. The UTD-learning algorithm (ϵ, δ) -learns C if for any target function $c^* \in C$, with probability $1 - \delta$, the algorithm outputs some hypothesis $h \in \mathcal{H}$, such that $\text{Err}_D(h, c^*) \leq v + \epsilon$.

3. Definitions And Notations

For a function class \mathcal{H} with domain \mathcal{Z} and range $\mathcal{Y} = \{1, \dots, l\}$, denote the zero-one loss class

$$L_{\mathcal{H}} = \{Z \times \{1, \dots, l\} \ni (z, y) \mapsto I_{h(z) \neq y} : h \in \mathcal{H}\}$$

For \mathcal{H} with domain \mathcal{Z} and range \mathbb{R} , denote the L_p loss class

$$L_{\mathcal{H}}^p = \{Z \times \mathbb{R} \ni (z, y) \mapsto |h(z) - y|^p : h \in \mathcal{H}\}$$

Remark. Throughout the article, we assume a bounded loss $\text{loss}(h(z), y) \leq M$. Without the loss of generality we use $M = 1$, otherwise, we can rescale M .

Define the following operations on the loss class $L_{\mathcal{H}}$. The convex hull of $L_{\mathcal{H}}$ is the set of all convex combinations of hypotheses from $L_{\mathcal{H}}$:

$$\text{conv}(L_{\mathcal{H}}) = \left\{ Z \times \mathcal{Y} \ni (z, y) \mapsto \sum_{t=1}^T \alpha_t f_t(z, y) : T \in \mathbb{N}, \alpha_t \in [0, 1], \sum_t \alpha_t = 1, f_t \in L_{\mathcal{H}} \right\}.$$

The convex hull of $L_{\mathcal{H}}$, where the data is corrupted by $\rho(\cdot)$, is denoted by

$$\text{conv}^{\rho}(L_{\mathcal{H}}) = \left\{ \mathcal{X} \times \mathcal{Y} \ni (x, y) \mapsto \max_{z \in \rho(x)} \sum_{t=1}^T \alpha_t f_t(z, y) : T \in \mathbb{N}, \alpha_t \in [0, 1], \sum_t \alpha_t = 1, f_t \in L_{\mathcal{H}} \right\}.$$

For $1 \leq j \leq k$ define,

$$\mathcal{F}_{\mathcal{H}}^{(j)} = \left\{ \mathcal{X} \times \mathcal{Y} \ni (x, y) \mapsto I_{\{h(z_j) \neq y\}} : h \in \mathcal{H}, \rho(x) = \{z_1, \dots, z_k\} \right\},$$

where we treat the set-valued output of $\rho(x)$ as an ordered list, and $\mathcal{F}_{\mathcal{H}}^{(j)}$ is constructed by taking the j th element in this list, for each input x .

For set W and k function classes $\mathcal{A}^{(1)}, \dots, \mathcal{A}^{(k)} \subseteq \mathbb{R}^W$, define the max operator

$$\max((\mathcal{A}^{(j)})_{j \in [k]}) := \left\{ W \ni w \mapsto \max_{j \in [k]} f^{(j)}(w) : f^{(j)} \in \mathcal{A}^{(j)} \right\}.$$

We also define a hybrid max-conv operator:

$$\text{max-conv}((\mathcal{A}^{(j)})_{j \in [k]}) := \left\{ W \ni w \mapsto \max_{j \in [k]} \sum_{t=1}^T \alpha_t f_t^{(j)}(w) : T \in \mathbb{N}, \alpha_t \in [0, 1], \sum_t \alpha_t = 1, f_t^{(j)} \in \mathcal{A}^{(j)} \right\}.$$

Note that

$$\text{max-conv}((\mathcal{A}^{(j)})_{j \in [k]}) \subseteq \max(\text{conv}((\mathcal{A}^{(j)})_{j \in [k]})),$$

and the containment will generally be strict, since the former requires the same choice of convex coefficients for all $\mathcal{A}^{(j)}$'s, while the latter allows distinct ones.

Remark. Observe that

$$\text{conv}^\rho(L_{\mathcal{H}}) = \text{max-conv}((\mathcal{F}_{\mathcal{H}}^{(j)})_{j \in [k]})$$

is the loss class of $\text{conv}(\mathcal{H})$ under corruption of $\rho(\cdot)$. We use the notation $\text{max-conv}((\mathcal{F}_{\mathcal{H}}^{(j)})_{j \in [k]})$ and exploit its structural properties.

Denote the error (risk) of hypothesis $g : \mathcal{Z} \mapsto \mathcal{Y}$ under corruption of $\rho(\cdot)$ by

$$R(g) = \mathbb{E}_{x \sim D} [\max_{z \in \rho(x)} \text{loss}(g(z), y)],$$

and the empirical error on sample S under corruption of $\rho(\cdot)$ by

$$\widehat{R}(g) = \frac{1}{|S|} \sum_{(x, y) \in S} \max_{z \in \rho(x)} \text{loss}(g(z), y).$$

3.1 Combinatorial dimensions and capacity measures

Rademacher Complexity Let \mathcal{H} be of real valued function class on the domain space \mathcal{Z} . Define the empirical Rademacher complexity on a given sequence $\mathbf{z} = (z_1, \dots, z_n) \in \mathcal{Z}^n$:

$$R_n(\mathcal{H}|\mathbf{z}) = E_\sigma \sup_{h \in \mathcal{H}} \frac{1}{n} \sum_{i=1}^n \sigma_i h(z_i).$$

Fat-Shattering Dimension For $\mathcal{F} \subset \mathbb{R}^{\mathcal{X}}$ and $\gamma > 0$, we say that \mathcal{F} γ -shatters a set $S = \{x_1, \dots, x_m\} \subset \mathcal{X}$ if there exists an $r = (r_1, \dots, r_m) \in \mathbb{R}^m$ such that for each $b \in \{-1, 1\}^m$ there is a function $f_b \in \mathcal{F}$ such that

$$\forall i \in [m] : \begin{cases} f_b(x_i) \geq r_i + \gamma & \text{if } b_i = 1 \\ f_b(x_i) \leq r_i - \gamma & \text{if } b_i = -1 \end{cases} .$$

We refer to r as the *shift*. The γ -fat-shattering dimension $\text{fat}_\gamma(\mathcal{F})$ is the size of the largest γ -shattered set (possibly ∞).

Graph Dimension Let $\mathcal{H} \subseteq \mathcal{Y}^{\mathcal{X}}$ be a categorical function class such that $\mathcal{Y} = [l] = \{1, \dots, l\}$. Let $S \subseteq \mathcal{X}$. We say that \mathcal{H} G -shatters S if there exists an $f : S \mapsto \mathcal{Y}$ such that for every $T \subseteq S$ there is a $g \in \mathcal{H}$ such that

$$\forall x \in T, g(x) = f(x) \text{ and } \forall x \in S \setminus T, g(x) \neq f(x).$$

The graph dimension of \mathcal{H} , denoted $d_G(\mathcal{H})$, is the maximal cardinality of a set that is G -shattered by \mathcal{H} .

Natarajan Dimension Let $\mathcal{H} \subseteq \mathcal{Y}^{\mathcal{X}}$ be a categorical function class such that $\mathcal{Y} = [l] = \{1, \dots, l\}$. Let $S \subseteq \mathcal{X}$. We say that \mathcal{H} N -shatters S if there exist $f_1, f_2 : S \mapsto \mathcal{Y}$ such that for every $y \in S$ $f_1(y) \neq f_2(y)$, and for every $T \subseteq S$ there is a $g \in \mathcal{H}$ such that

$$\forall x \in T, g(x) = f_1(x), \text{ and } \forall x \in S \setminus T, g(x) = f_2(x).$$

The Natarajan dimension of \mathcal{H} , denoted $d_N(\mathcal{H})$, is the maximal cardinality of a set that is N -shattered by \mathcal{H} .

Growth Function The growth function $\Pi_{\mathcal{H}} : \mathbb{N} \mapsto \mathbb{N}$ for a binary function class $\mathcal{H} : \mathcal{X} \mapsto \{0, 1\}$ is defined by

$$\forall m \in \mathbb{N}, \Pi_{\mathcal{H}}(m) = \max_{\{x_1, \dots, x_m\} \subseteq \mathcal{X}} |\{(h(x_1), \dots, h(x_m)) : h \in \mathcal{H}\}|$$

And the VC-dimension of \mathcal{H} is defined by

$$\text{VC}(\mathcal{H}) = \max \{m : \Pi_{\mathcal{H}}(m) = 2^m\} .$$

4. Algorithm

We have a base hypothesis class \mathcal{H} with domain \mathcal{Z} and range \mathcal{Y} that can be $\{1, \dots, l\}$ or \mathbb{R} . The learner receives a labeled uncorrupted sample and has access during the training to possible corruptions by the adversary. We employ the regret minimization algorithm proposed by [Feige et al. \(2015\)](#) for binary classification, and extend it to the regression and multi-class classification settings.

A brief description of the algorithm is as follows. Given $x \in \mathcal{X}$, we define a $|\rho(x)| \times \mathcal{H}$ matrix M_x such that $M_x(z, h) = I_{\{h(z) \neq y\}}$, where $y = c^*(x)$. The learner's strategy is a distribution Q over \mathcal{H} . The adversary's strategy for a given $x \in \mathcal{X}$ is a distribution P_x over the corrupted inputs $\rho(x)$, we can treat P as a vector of probabilities P_x over all $x \in \mathcal{X}$. Via the minimax principle, the value of the game is

$$v = \min_Q \max_P \mathbb{E}_{x \sim D} [P_x^T M_x Q] = \max_P \min_Q \mathbb{E}_{x \sim D} [P_x^T M_x Q]$$

For a given P , a learner's minimizing Q is simply a hypothesis that minimizes the error when the distribution over pairs $(z, y) \in \mathcal{Z} \times \mathcal{Y}$ is D^P , where

$$D^P(z, y) = \sum_{x: c^*(x)=y \wedge z \in \rho(x)} P_x(z) D(x).$$

Hence, the learner selects

$$h^P = \arg \min_{h \in \mathcal{H}} \mathbb{E}_{(z,y) \sim D^P} [\text{loss}(h(z), y)].$$

A hypotheses h^P can be found using the ERM oracle, when D is the empirical distribution over a training sample.

Algorithm 1

parameter: $\eta > 0$

- 1: **for all** $(x, y) \in S, z \in \rho(x)$ **do** ▷ initialize weights and distributions vector
- 2: $w_1(z, (x, y)) \leftarrow 1, \forall (x, y) \in S, \forall z \in \rho(x)$
- 3: $P^1(z, (x, y)) \leftarrow \frac{w_1(z, (x, y))}{\sum_{z' \in \rho(x)} w_1(z', (x, y))}$ ▷ for each $(x, y) \in S$ we have a distribution
- 4: **end for**
- 5: **for** $t = 1:T$ **do**
- 6: $h_t \leftarrow \arg \min_{h \in \mathcal{H}} E_{(z,y) \sim D^{P^t}} [\text{loss}(h(z), y)]$ ▷ using the ERM oracle for H
- 7: **for all** $(x, y) \in S, z \in \rho(x)$ **do** ▷ update weights and distributions vector
- 8: $w_{t+1}(z, (x, y)) \leftarrow (1 + \eta \cdot [\text{loss}(h_t(z), y)] \cdot w_t(z, (x, y)))$
- 9: $P^{t+1}(z, (x, y)) \leftarrow \frac{w_{t+1}(z, (x, y))}{\sum_{z' \in \rho(x)} w_{t+1}(z', (x, y))}$
- 10: **end for**
- 11: **end for**
- 12: **return** $Q = \frac{1}{T} \sum_{t=1}^T h_t, P = \frac{1}{T} \sum_{t=1}^T P^t$ ▷ Q for the learner, P for the adversary

Similar to Feige et al. (2015, Theorem 1), for the binary classification case and zero-one loss we have:

Theorem 1 (Feige, Mansour, and Schapire, 2015, Theorem 1) *Fix a sample S , and let $T \geq \frac{4|S|\ln(m)}{\epsilon^2}$. For an uncorrupted input domain S we have that the strategy $P = \frac{1}{T} \sum_{t=1}^T P^t$ for the adversary is an ϵ -optimal strategy and the strategy $Q = \frac{1}{T} \sum_{t=1}^T h_t$ for the learner is an ϵ -optimal strategy.*

Assuming a bounded loss, i.e., $\text{loss}(h(z), y) \leq 1, \forall x \in \mathcal{X}, \forall z \in \mathcal{Z}, \forall h \in \mathcal{H}$, the result remains the same for the other settings.

5. Generalization Bound For Binary Classification

We would like to show that if the sample S is large enough, then the policy achieved by the algorithm above will generalize well. We both improve a generalization bound, previously found in Feige et al. (2015), which handles a weighted average of hypotheses from \mathcal{H} , and also are able to handle an infinite hypothesis class \mathcal{H} . The sample complexity is improved from $\mathcal{O}(\frac{1}{\epsilon^4} \log(\frac{|\mathcal{H}|}{\delta}))$ to $\mathcal{O}(\frac{1}{\epsilon^2} (k \log(k) \text{VC}(\mathcal{H}) + \log \frac{1}{\delta}))$.

Theorem 2 Let $\mathcal{H} : \mathcal{Z} \mapsto \{0, 1\}$ be a hypothesis class with finite VC-dimension. There is a sample complexity $m_0 = \mathcal{O}(\frac{1}{\epsilon^2}(k \log(k) \text{VC}(\mathcal{H}) + \log \frac{1}{\delta}))$, such that for $|S| \geq m_0$, for every $g \in \text{conv}(\mathcal{H})$,

$$|R(g) - \widehat{R}(g)| \leq \epsilon$$

with probability at least $1 - \delta$.

Lemma 3 For any k real valued function classes $\mathcal{F}^{(1)}, \dots, \mathcal{F}^{(k)}$, over a set \mathcal{X} and $\mathbf{x} = (x_1, \dots, x_n) \in \mathcal{X}^n$, we have

$$R_n(\text{max-conv}((\mathcal{F}^{(j)})_{j \in [k]} | \mathbf{x})) = R_n(\text{max}((\mathcal{F}^{(j)})_{j \in [k]} | \mathbf{x})).$$

Proof It is easily seen that $\text{max}((\mathcal{F}^{(j)})_{j \in [k]} | \mathbf{x}) \subseteq \text{max-conv}((\mathcal{F}^{(j)})_{j \in [k]} | \mathbf{x})$, since for any $f^{(j)} \in \mathcal{F}^{(j)}$, $j \in [k]$ and $T = 1$ we have that $\max_j f^{(j)} \in \text{max-conv}((\mathcal{F}^{(j)})_{j \in [k]} | \mathbf{x})$. This proves that the right-hand side is at least as large as the left-hand side. Conversely,

$$\begin{aligned} R_n(\text{max-conv}((\mathcal{F}^{(j)})_{j \in [k]} | \mathbf{x})) &= E_\sigma \sup_{\substack{(f_t^{(j)})_{j \in [k]}, t \in [T] \\ (\alpha_t)_{t \in [T]}} \frac{1}{n} \sum_{i=1}^n \sigma_i \max_{j \in [k]} \sum_{t=1}^T \alpha_t f_t^{(j)}(x_i) \\ &\leq E_\sigma \sup_{\substack{(f_t^{(j)})_{j \in [k]}, t \in [T] \\ (\alpha_t)_{t \in [T]}} \frac{1}{n} \sum_{i=1}^n \sigma_i \sum_{t=1}^T \alpha_t \max_{j \in [k]} f_t^{(j)}(x_i) \\ &= R_n(\text{conv}(\text{max}((\mathcal{F}^{(j)})_{j \in [k]} | \mathbf{x})) = R_n(\text{max}((\mathcal{F}^{(j)})_{j \in [k]} | \mathbf{x})), \end{aligned}$$

where the last equality stems from the well-known identity $R_n(\mathcal{F} | \mathbf{x}) = R_n(\text{conv}(\mathcal{F}) | \mathbf{x})$ (Boucheron et al., 2005, Theorem 3.3). ■

Lemma 4 Let $\Psi : \mathcal{F}_1 \times \dots \times \mathcal{F}_k \rightarrow \mathcal{F}$ be an arbitrary mapping, where $\mathcal{F}, \mathcal{F}_1, \dots, \mathcal{F}_k \subseteq \{-1, 1\}^X$ and $\text{VC}(\mathcal{F}_j) = d_j$ for $j \in [k]$. Then the VC-dimension of $\Psi(\mathcal{F}_1 \times \dots \times \mathcal{F}_k)$ is less than $2k \log(3k) \bar{d}$, where $\bar{d} := \frac{1}{k} \sum_{i=1}^k d_j$.

Proof We adapt the argument of Blumer et al. (1989, Lemma 3.2.3), which is stated therein for k -fold unions and intersections. The $k = 1$ case is trivial, so assume $k \geq 2$. For any $S \subseteq \mathcal{X}$, define $\Psi(\mathcal{F}_1 \times \dots \times \mathcal{F}_k)(S) \subseteq \{-1, 1\}^S$ to be the restriction of $\Psi(\mathcal{F}_1 \times \dots \times \mathcal{F}_k)$ to S . The key observation is that

$$\begin{aligned} |\Psi(\mathcal{F}_1 \times \dots \times \mathcal{F}_k)(S)| &\leq \prod_{j=1}^k |\mathcal{F}_j(S)| \\ &\leq \prod_{j=1}^k (e|S|/d_j)^{d_j} \\ &\leq (e|S|/\bar{d})^{\bar{d}k}. \end{aligned}$$

The last inequality requires proof. After taking logarithms and dividing both sides by k , it is equivalent to the claim that

$$\bar{d} \log \bar{d} \leq \frac{1}{k} \sum_{j=1}^k d_j \log d_j$$

— an immediate consequence of Jensen’s inequality applied to the convex function $f(x) = x \log x$.

The rest of the argument is identical that of [Blumer et al.](#): one readily verifies that for $m = |S| = 2\bar{d}k \log(3k)$, we have $(em/\bar{d})^{\bar{d}k} < 2^m$. ■

Theorem 5 ([Mohri et al., 2012, Theorem 3.1](#)) *Let \mathcal{F} be a family of functions mapping from \mathcal{Z} to $[0, 1]$. Denote $E(f) := E_{z \sim D}[f(z)]$ and $\hat{E}_S(f) = \frac{1}{n} \sum_{i=1}^n f(z_i)$. Then, for every $\delta > 0$ with probability at least $1 - \delta$, for all $f \in \mathcal{F}$:*

$$E[f] - \hat{E}_S(f) \leq 2R_n(\mathcal{F}|\mathbf{z}) + 3\sqrt{\frac{\ln(2/\delta)}{2n}}$$

Theorem 6 ([Dudley, 1967; Bartlett and Mendelson, 2002](#)) *Let $d = \text{VC}(\mathcal{F})$, then for $m \geq d \geq 1$ and for some absolute constant $c > 0$:*

$$R_n(\mathcal{F}|\mathbf{z}) \leq c\sqrt{\frac{d}{|S|}}.$$

Proof of Theorem 2 Our strategy is to bound the empirical Rademacher complexity of the loss class of $\text{conv}(\mathcal{H})$. As we mentioned in Section 3, the loss class is $\text{conv}^\rho(L_{\mathcal{H}}) = \text{max-conv}((\mathcal{F}_{\mathcal{H}}^{(j)})_{j \in [k]})$, we use the notation $\text{max-conv}((\mathcal{F}_{\mathcal{H}}^{(j)})_{j \in [k]})$. Recall that functions contained in $\mathcal{F}_{\mathcal{H}}^{(j)}$ are loss functions of the learner when the adversary corrupts input x to $z_j \in \rho(x)$. Combine everything together,

$$\begin{aligned} |R(g) - \hat{R}(g)| &= |E_{(x,y) \sim D} \max_{j \in [k]} \sum_{t=1}^T \alpha_t f_t^{(j)}(x) - \frac{1}{|S|} \sum_{(x,y) \in S} \max_{j \in [k]} \sum_{t=1}^T \alpha_t f_t^{(j)}(x)| \\ &\stackrel{(i)}{\leq} 2R_n(\text{max-conv}((\mathcal{F}_{\mathcal{H}}^{(j)})_{j \in [k]})|\mathbf{z}) + 3\sqrt{\frac{\ln(2/\delta)}{2|S|}} \\ &\stackrel{(ii)}{=} 2R_n(\text{max}((\mathcal{F}_{\mathcal{H}}^{(j)})_{j \in [k]})|\mathbf{z}) + 3\sqrt{\frac{\ln(2/\delta)}{2|S|}} \\ &\stackrel{(iii)}{\leq} 2c\sqrt{\frac{\text{VC}(\text{max}((\mathcal{F}_{\mathcal{H}}^{(j)})_{j \in [k]}))}{|S|}} + 3\sqrt{\frac{\ln(2/\delta)}{2|S|}} \\ &\stackrel{(iv)}{\leq} 2c\sqrt{\frac{2k \log(3k) \max_{j \in [k]} \text{VC}(\mathcal{F}_{\mathcal{H}}^{(j)})}{|S|}} + 3\sqrt{\frac{\ln(2/\delta)}{2|S|}} \\ &\stackrel{(v)}{\leq} 2c\sqrt{\frac{2k \log 3k \text{VC}(\mathcal{H})}{|S|}} + 3\sqrt{\frac{\ln(2/\delta)}{2|S|}} \\ &\leq \epsilon, \end{aligned}$$

where (i) stems from Theorem 5 (generalization error by Rademacher complexity bound for the function class $\max\text{-conv}((\mathcal{F}_{\mathcal{H}}^{(j)})_{j \in [k]})$, (ii) stems from Lemma 3, (iii) stems from Theorem 6 (bound on Rademacher complexity using VC-dimension), (iv) stems from Lemma 4 and (v) stems from Lemma 8. ■

6. Generalization Bound For Multi-Class Classification

Let $\mathcal{H} \subseteq \mathcal{Y}^{\mathcal{Z}}$ be a function class such that $\mathcal{Y} = [l] = \{1 \dots, l\}$. We follow similar arguments to the binary case.

Theorem 7 *Let \mathcal{H} be a function class with domain \mathcal{Z} and range $\mathcal{Y} = [l]$ with finite Graph-dimension $d_G(\mathcal{H})$. There is a sample complexity $m_0 = \mathcal{O}(\frac{1}{\epsilon^2}(k \log(k)d_G(\mathcal{H}) + \log \frac{1}{\delta}))$, such that for $|S| \geq m_0$, for every $g \in \text{conv}(\mathcal{H})$,*

$$|R(g) - \widehat{R}(g)| \leq \epsilon$$

with probability at least $1 - \delta$.

Lemma 8 *Let \mathcal{H} be a function class with domain \mathcal{Z} and range $\mathcal{Y} = [l]$. Denote the Graph-dimension of \mathcal{H} by $d_G(\mathcal{H})$. Then*

$$\text{VC}(\mathcal{F}_{\mathcal{H}}^{(j)}) \leq d_G(\mathcal{H}).$$

In particular, for binary-valued classes, $\text{VC}(\mathcal{F}_{\mathcal{H}}^{(j)}) \leq \text{VC}(\mathcal{H})$ — since for these, the VC- and Graph-dimensions coincide.

For the proof of Theorem 7, we follow the same proof of Theorem 2 and use the Graph-dimension property of Lemma 8 in the (v) inequality.

Remark. A similar bound to that of Theorem 2 can be achieved by using the Natarajan dimension and the fact that

$$d_G(\mathcal{H}) \leq 4.67 \log_2(|\mathcal{Y}|)d_N(\mathcal{H})$$

as previously shown [Ben-David et al. \(1995\)](#).

7. Generalization Bound For Regression

Let $\mathcal{H} \subseteq \mathbb{R}^{\mathcal{Z}}$ be a hypothesis class of real functions. We refer to a bounded regression problem, we assume a bounded loss function by 1.

In order to use similar arguments to the binary case, we need an analogous to Lemma 4 for the fat-shattering dimension and understand the connection between the shattering dimension of loss classes (L_1 and L_2) to the original function class.

Theorem 9 *Let \mathcal{H} be a function class with domain \mathcal{Z} and range \mathbb{R} . Assume \mathcal{H} has a finite γ -fat-shattering dimension for all $\gamma > 0$. Denote $m_{\mathcal{H}}(\gamma) = \int_0^1 \sqrt{\text{fat}_{c\gamma}(\mathcal{H}) \log(\frac{2}{\gamma})} d\gamma$, where $c > 0$ is an*

absolute constant. For the ℓ_1 loss function, there is a sample complexity $m_0 = \mathcal{O}(\frac{1}{\epsilon^2}(k \log(k)m_{\mathcal{H}}(\gamma) + \log \frac{1}{\delta}))$, such that for $|S| \geq m_0$, for every $g \in \text{conv}(\mathcal{H})$,

$$|R(g) - \widehat{R}(g)| \leq \epsilon$$

with probability at least $1 - \delta$.

Remark In case of integral divergence we can use a refined version of Dudley’s entropy integral as in Theorem 18.

Corollary 10 Let \mathcal{H} be a function class of homogeneous hyperplanes with domain \mathbb{R}^m . Using the same assumptions as in Theorem 9, the sample complexity is $m_0 = \mathcal{O}(\frac{1}{\epsilon^2}(k \log^2(k/\epsilon) + \log \frac{1}{\delta}))$.

Corollary 11 For the ℓ_2 loss the same result of Theorem 9 holds when we redefine $m_{\mathcal{H}}(\gamma) = \int_0^1 \sqrt{\text{fat}_{c\gamma/2}(\mathcal{H}) \log(\frac{2}{\gamma})} d\gamma$.

7.1 Shattering dimension of the class $\max((\mathcal{A}^{(j)})_{j \in [k]})$

The main result of this section is bounding the fat shattering dimension of $\max((\mathcal{A}^{(j)})_{j \in [k]})$ class.

Theorem 12 For any k real valued functions classes $\mathcal{F}_1, \dots, \mathcal{F}_k$ with finite fat-shattering dimension, we have

$$\text{fat}_{\gamma}(\max((\mathcal{F}_j)_{j \in [k]})) < 2 \log(3k) \sum_{j=1}^k \text{fat}_{\gamma}(\mathcal{F}_j)$$

for all $\gamma > 0$.

Remark. This result generalizes an analogous bound obtained in Kontorovich (2018) for maxima of linear classes. For a fixed scale $\gamma > 0$, Csikos et al. (2018) have recently shown that the $O(\log k)$ factor cannot, in general, be removed. Whether a single function class can attain the lower bound for every $\gamma > 0$ simultaneously is an open problem.

We begin with an auxiliary definition. We say that \mathcal{F} “ γ -shatters a set S at zero” if the shift (or witness) r is constrained to be 0 in the the usual γ -shattering definition (has appeared previously in Gottlieb et al. (2014)). The analogous dimension will be denoted by $\text{fat}_{\gamma}^0(\mathcal{F})$.

Lemma 13 For all $\mathcal{F} \subseteq \mathbb{R}^{\mathcal{X}}$ and $\gamma > 0$, we have

$$\text{fat}_{\gamma}(\mathcal{F}) = \max_{r \in \mathbb{R}^{\mathcal{X}}} \text{fat}_{\gamma}^0(\mathcal{F} - r), \tag{1}$$

where $\mathcal{F} - r = \{f - r : f \in \mathcal{F}\}$ is the r -shifted class; in particular, the maximum is always achieved.

Proof Fix \mathcal{F} and γ . For any choice of $r \in \mathbb{R}^{\mathcal{X}}$, if $\mathcal{F} - r$ γ -shatters some set $S \subseteq \mathcal{X}$ at zero, then then \mathcal{F} γ -shatters S in the usual sense with shift $r_S \in \mathbb{R}^S$ (i.e., the restriction of r to S). This proves that the left-hand side of (1) is at least as large as the right-hand side. Conversely, suppose that \mathcal{F} γ -shatters some $S \subseteq \mathcal{X}$ in the usual sense, with some shift $r \in \mathbb{R}^S$. Choosing $r' \in \mathbb{R}^{\mathcal{X}}$ by $r'_S = r$ and $r'_{\mathcal{X} \setminus S} = 0$, we see that $\mathcal{F} - r'$ γ -shatters S at zero. This proves the other direction and hence the claim. ■

Lemma 14 *Suppose that $F \subseteq \{-1, 1, \star\}^m$ has VC-dimension d in the sense that some $J \subseteq [m]$ of size d verifies $F(J) = \{-1, 1\}^J$ and for all $J' \subseteq [m]$ with $|J'| > d$, we have $F(J') \subsetneq \{-1, 1\}^{J'}$. Then there is a mapping $\varphi : F \rightarrow \{-1, 1\}^m$ such that (i) for all $v \in F$ and all $i \in [m]$, we have $v_i \neq \star \implies (\varphi(v))_i = v_i$ and (ii) $\varphi(F)$ does not shatter more than d points.*

Proof The mapping φ must resolve each ‘‘ambiguity’’ $v_i = \star$ as $(\varphi(v))_i \in \{-1, 1\}$ in such a way that the resulting set of vectors $\varphi(F)$ does not shatter more points than F does. We achieve this via an iterative procedure, which initializes $F' := F$ and modifies each $v \in F'$, element-wise, until $F' \subseteq \{-1, 1\}^m$ — that is, all of the ambiguities have been resolved.

Suppose that the VC-dimension of F' is d and some $v \in F'$ and $i \in [m]$ are such that $v_i = \star$; we must choose a value for $(\varphi(v))_i \in \{-1, 1\}$. If one of these choices ensures the condition that the VC-dimension will not increase, then we’re done. Otherwise, the VC-dimension will increase from d to $d + 1$ for *both* choices of $(\varphi(v))_i = 1$ and $(\varphi(v))_i = -1$. This means, in particular, that F shatters some set $J \subseteq [m]$ of size d and $i \notin J$ — since otherwise, disambiguating v_i from \star to ± 1 would not increase the VC-dimension. Since the choice $(\varphi(v))_i = 1$ increases the VC-dimension, it must be the case that

$$F(J \cup \{i\}) = \{-1, 1\}^{J \cup \{i\}} \setminus \{z_{J \cup \{i\}}\} \quad (2)$$

for some ‘‘missing witness’’ $z \in \{-1, 1, \star\}^m$, which agrees with v on J and $z_i = 1$; the notation z_E indicates the restriction of Z to the index set $E \subseteq [m]$. Analogously, since the choice $(\varphi(v))_i = -1$ also increases the VC-dimension, we have

$$F(J \cup \{i\}) = \{-1, 1\}^{J \cup \{i\}} \setminus \{z'_{J \cup \{i\}}\}, \quad (3)$$

where $z'_j = v_j$ and $z'_i = -1$. The conditions (2) and (3) are in obvious contradiction, from which we conclude that the ambiguity can be resolved for each $v_i = \star$ without increasing the VC-dimension. ■

Proof of Theorem 12 First, we observe that r -shift commutes with the max operator:

$$\max((\mathcal{F}_j - r)_{j \in [k]}) = \max((\mathcal{F}_j)_{j \in [k]}) - r$$

and so, in light of Lemma 13, we have

$$\text{fat}_\gamma(\max((\mathcal{F}_j)_{j \in [k]})) = \max_r \text{fat}_\gamma^0(\max((\mathcal{F}_j)_{j \in [k]}) - r) = \max_r \text{fat}_\gamma^0(\max((\mathcal{F}_j - r)_{j \in [k]})).$$

Hence, to prove Theorem 12, it suffices to show that

$$\text{fat}_\gamma^0(\max((\mathcal{F}_j)_{j \in [k]})) \leq 2 \log(3k) \sum_{j=1}^k \text{fat}_\gamma^0(\mathcal{F}_j). \quad (4)$$

To prove (4), let us fix some $S = \{x_1, \dots, x_m\} \subset \mathcal{X}$ and convert each $\mathcal{F}_j(S) \subseteq \mathbb{R}^m$ to a finite class $\mathcal{F}_j^*(S) \subseteq \{-\gamma, \gamma, \star\}^m$ as follows. For every vector in $v \in \mathcal{F}_j(S)$, define $v^* \in \mathcal{F}_j^*(S)$ by: $v_i^* = \text{sgn}(v_i)\gamma$ if $|v_i| \geq \gamma$ and $v_i^* = \star$ else. The notion of shattering (at zero) remains the same: a set $T \subseteq S$ is shattered by \mathcal{F}_j iff $\mathcal{F}_j^*(T) = \{-\gamma, \gamma\}^T$. Lemma 14 furnishes a mapping $\varphi : \mathcal{F}_j^*(S) \rightarrow \{-\gamma, \gamma\}^m$ such that (i) for all $v \in \mathcal{F}_j^*(S)$ and all $i \in [m]$, we have $v_i \neq \star \implies (\varphi(v))_i = v_i$ and (ii)

$\varphi(\mathcal{F}_j^*(S))$ does not shatter more points than $\mathcal{F}_j^*(S)$. Together, properties (i) and (ii) imply that $\text{fat}_\gamma^0(\mathcal{F}_j(S)) = \text{fat}_\gamma^0(\varphi(\mathcal{F}_j^*(S)))$ for all j .

Finally, observe that any D points in S γ -shattered by $\max(\mathcal{F}_{j \in [k]})$ are also shattered by $\max(\varphi(\mathcal{F}_{j \in [k]}^*(S)))$. Applying Lemma 4 with $\Psi(f_1, \dots, f_k)(x) = \max_{j \in [k]} f_j(x)$ shows that $\max(\varphi(\mathcal{F}_{j \in [k]}^*(S)))$ cannot shatter $2 \log(3k) \sum_{j=1}^k d_j$ points, where $d_j = \text{fat}_\gamma^0(\varphi(\mathcal{F}_j^*(S))) = \text{fat}_\gamma^0(\mathcal{F}_j(S)) \leq \text{fat}_\gamma^0(\mathcal{F}_j)$. We have shown that, for all finite $S \subseteq \mathcal{X}$, we have $\text{fat}_\gamma^0(\max(\mathcal{F}_{j \in [k]}(S))) \leq 2 \log(3k) \sum_{j=1}^k \text{fat}_\gamma^0(\mathcal{F}_j(S))$.

Since this latter estimate holds independently of S , it is also an upper bound on $\text{fat}_\gamma^0(\max(\mathcal{F}_{j \in [k]}))$. ■

7.2 Shattering dimension of L_1 and L_2 loss classes

Lemma 15 *Let $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$ be a real valued function class. denote $L_{\mathcal{H}}^1$ and $L_{\mathcal{H}}^2$ the L_1 and L_2 loss classes of \mathcal{H} respectively. Assume $L_{\mathcal{H}}^1$ is bounded by M . For any \mathcal{H} ,*

$$\text{fat}_\gamma(L_{\mathcal{H}}^1) \leq 2 \text{fat}_\gamma(\mathcal{H}) + 1, \text{fat}_\gamma(L_{\mathcal{H}}^2) \leq 2 \text{fat}_{\gamma/2M}(\mathcal{H}) + 1.$$

Lemma 16 *For $p \in \{1, 2\}$ and $j \in [k]$, define*

$$\mathcal{F}_{\mathcal{H}}^{(p,j)} = \{\mathcal{X} \times \mathcal{Y} \ni (x, y) \mapsto |h(z_j) - y|^p : h \in \mathcal{H}, \rho(x) = \{z_1, \dots, z_k\}\}.$$

Then

$$\text{conv}^\rho(L_{\mathcal{H}}^p) = \max\text{-conv}((\mathcal{F}_{\mathcal{H}}^{(p,j)})_{j \in [k]})$$

and, for all $\gamma > 0$,

$$\text{fat}_\gamma(\mathcal{F}_{\mathcal{H}}^{(p,j)}) \leq \text{fat}_\gamma(L_{\mathcal{H}}^p).$$

Proof The first claim is immediate from the definitions, while the second is proved using the argument (almost verbatim) of Lemma 8. ■

Lemma 17 *For $\mathcal{H}, \mathcal{G} \subseteq \{-1, 1\}^{\mathcal{X}}$ with finite VC-dimension, we have*

$$\text{VC}(\mathcal{H} \cup \mathcal{G}) \leq \text{VC}(\mathcal{H}) + \text{VC}(\mathcal{G}) + 1,$$

and for $\mathcal{H}, \mathcal{G} \subseteq \mathbb{R}^{\mathcal{X}}$ with finite γ -fat-shattering dimension, we have

$$\text{fat}_\gamma(\mathcal{H} \cup \mathcal{G}) \leq \text{fat}_\gamma(\mathcal{H}) + \text{fat}_\gamma(\mathcal{G}) + 1.$$

Proof [of Lemma 15] For any \mathcal{X} and any function class $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$, define the *difference class* $\mathcal{H}^\Delta \subset \mathbb{R}^{\mathcal{X} \times \mathbb{R}}$ as

$$\mathcal{H}^\Delta = \{\mathcal{X} \times \mathbb{R} \ni (x, y) \mapsto \Delta_h(x, y) := h(x) - y; h \in \mathcal{H}\}.$$

In words: \mathcal{H}^Δ consists of all functions $\Delta_h(x, y) = h(x) - y$ indexed by $h \in \mathcal{H}$.

It is easy to see that for all $\gamma > 0$, we have $\text{fat}_\gamma(\mathcal{H}^\Delta) \leq \text{fat}_\gamma(\mathcal{H})$. Indeed, if \mathcal{H}^Δ γ -shatters some set $\{(x_1, y_1), \dots, (x_k, y_k)\} \subset \mathcal{X} \times \mathbb{R}$ with shift $r \in \mathbb{R}^k$, then \mathcal{H} γ -shatters the set $\{x_1, \dots, x_k\} \subset \mathcal{X}$ with shift $r + (y_1, \dots, y_k)$.

Now let $\mathcal{F} \subset [0, M]^\mathcal{X}$ and define $\mathcal{F}^{\circ 2} := \{f^2; f \in \mathcal{F}\}$. We would like to bound $\text{fat}_\gamma(\mathcal{F}^{\circ 2})$ in terms of $\text{fat}_\gamma(\mathcal{F})$. Suppose that $\mathcal{F}^{\circ 2}$ γ -shatters some set $\{x_1, \dots, x_k\}$ with shift $r^2 = (r_1^2, \dots, r_k^2) \in [0, M]^k$ (there is no loss of generality in assuming that the shift has the same range as the function class). Using the elementary inequality

$$|a^2 - b^2| \leq 2M|a - b|, \quad a, b \in [0, M],$$

we conclude that \mathcal{F} is able to $\gamma/(2M)$ -shatter the same k points and thus $\text{fat}_\gamma(\mathcal{F}^{\circ 2}) \leq \text{fat}_{\gamma/(2M)}(\mathcal{F})$.

To generalize this result to the case where $\mathcal{F} \subset [-M, M]^\mathcal{X}$, we decompose \mathcal{F} as $\mathcal{F} = \mathcal{F}_+ \cup \mathcal{F}_-$, where $\mathcal{F}_+ = \mathcal{F} \cap [0, M]^\mathcal{X}$ and $\mathcal{F}_- = \mathcal{F} \cap [-M, 0]^\mathcal{X}$. Clearly, then, $\mathcal{F}^{\circ 2} = (\mathcal{F}_+)^{\circ 2} \cup (\mathcal{F}_-)^{\circ 2}$. Furthermore, using Lemma 17 we observe that for any $\mathcal{H}, \mathcal{G} \subset \mathbb{R}^\mathcal{X}$, we have

$$\text{fat}_\gamma(\mathcal{H} \cup \mathcal{G}) \leq \text{fat}_\gamma(\mathcal{H}) + \text{fat}_\gamma(\mathcal{G}) + 1,$$

whence

$$\begin{aligned} \text{fat}_\gamma(\mathcal{F}^{\circ 2}) &\leq \text{fat}_\gamma((\mathcal{F}_+)^{\circ 2}) + \text{fat}_\gamma((\mathcal{F}_-)^{\circ 2}) + 1 \\ &\leq \text{fat}_{\gamma/(2M)}(\mathcal{F}_+) + \text{fat}_{\gamma/(2M)}(\mathcal{F}_-) + 1 \\ &\leq 2 \text{fat}_{\gamma/(2M)}(\mathcal{F}) + 1, \end{aligned}$$

where the second inequality follows from the fact that $\text{fat}_\gamma(-\mathcal{F}) = \text{fat}_\gamma(\mathcal{F})$.

Finally, define \mathcal{F} as the ℓ_2 loss class of \mathcal{H} :

$$\mathcal{F} = \{\mathcal{X} \times \mathbb{R} \ni (x, y) \mapsto (h(x) - y)^2; h \in \mathcal{H}\}.$$

Then

$$\begin{aligned} \text{fat}_\gamma(\mathcal{F}) &= \text{fat}_\gamma((\mathcal{H}^\Delta)^{\circ 2}) \\ &\leq 2 \text{fat}_{\gamma/(2M)}(\mathcal{H}^\Delta) + 1 \\ &\leq 2 \text{fat}_{\gamma/(2M)}(\mathcal{H}) + 1. \end{aligned}$$

The case of ℓ_1 is similar, $\mathcal{F} \subset [0, M]^\mathcal{X}$ define $\text{abs}(\mathcal{F}) := \{|f|; f \in \mathcal{F}\}$. Clearly, $\text{fat}_\gamma(\text{abs}(\mathcal{F})) = \text{fat}_\gamma(\mathcal{F})$. Because \mathcal{F} is positive. Then,

$$\begin{aligned} \text{fat}_\gamma(\text{abs}(\mathcal{F})) &\leq \text{fat}_\gamma(\text{abs}(\mathcal{F}_+)) + \text{fat}_\gamma(\text{abs}(\mathcal{F}_-)) + 1 \\ &= 2 \text{fat}_\gamma(\mathcal{F}) + 1, \end{aligned}$$

Finally, define \mathcal{F} as the ℓ_1 loss class of \mathcal{H} :

$$\mathcal{F} = \{\mathcal{X} \times \mathbb{R} \ni (x, y) \mapsto |h(x) - y|; h \in \mathcal{H}\}.$$

Then

$$\begin{aligned} \text{fat}_\gamma(\mathcal{F}) &= \text{fat}_\gamma(\text{abs}(\mathcal{H}^\Delta)) \\ &\leq 2 \text{fat}_\gamma(\mathcal{H}^\Delta) + 1 \\ &\leq 2 \text{fat}_\gamma(\mathcal{H}) + 1. \end{aligned}$$

■

7.3 Generalization bound proof

Theorem 18 (*Dudley, 1967; Mendelson and Vershynin, 2003*) For any $\mathcal{F} \subseteq [-1, 1]^X$ and any $\gamma \in (0, 1)$,

$$R_n(\mathcal{F}|\mathbf{z}) \leq 12\sqrt{\frac{\tilde{K}}{n}} \int_0^1 \sqrt{\text{fat}_{c\gamma}(\mathcal{F}) \log\left(\frac{2}{\gamma}\right)} d\gamma$$

where c and \tilde{K} are universal constants.

Remark In case of integral divergence we can use a refined version of Dudley's entropy integral:

$$R_n(\mathcal{F}|\mathbf{z}) \leq \inf_{\alpha \geq 0} \left\{ 4\alpha + 12\sqrt{\frac{\tilde{K}}{n}} \int_{\alpha}^1 \sqrt{\text{fat}_{c\gamma}(\mathcal{F}) \log\left(\frac{2}{\gamma}\right)} d\gamma \right\}$$

Proof [of Theorem 9] Similar to the proof for binary case, we bound the empirical Rademacher complexity of the loss class of $\text{conv}(\mathcal{H})$.

$$\begin{aligned} |R(g) - \widehat{R}(g)| &= |E_{(x,y) \sim D} \max_{j \in [k]} \sum_{t=1}^T \alpha_t f_t^{(j)}(x) - \frac{1}{|S|} \sum_{(x,y) \in S} \max_{j \in [k]} \sum_{t=1}^T \alpha_t f_t^{(j)}(x)| \\ &\stackrel{(i)}{\leq} 2R_n(\text{max-conv}((\mathcal{F}_{\mathcal{H}}^{(j)})_{j \in [k]})|\mathbf{z}) + 3\sqrt{\frac{\ln(2/\delta)}{2|S|}} \\ &\stackrel{(ii)}{\leq} 2R_n(\text{max}((\mathcal{F}_{\mathcal{H}}^{(j)})_{j \in [k]})) + 3\sqrt{\frac{\ln(2/\delta)}{2|S|}} \\ &\stackrel{(iii)}{\leq} 24\sqrt{\frac{\tilde{K}}{|S|}} \int_0^1 \sqrt{\text{fat}_{c\gamma}(\text{max}((\mathcal{F}_{\mathcal{H}}^{(j)})_{j \in [k]})) \log\left(\frac{2}{\gamma}\right)} d\gamma + 3\sqrt{\frac{\ln(2/\delta)}{2|S|}} \\ &\stackrel{(iv)}{\leq} 24\sqrt{\frac{2k \log(3k) \tilde{K}}{|S|}} \int_0^1 \sqrt{\max_{j \in [k]} \text{fat}_{c\gamma}(\mathcal{F}_{\mathcal{H}}^{(j)}) \log\left(\frac{2}{\gamma}\right)} d\gamma + 3\sqrt{\frac{\ln(2/\delta)}{2|S|}} \\ &\stackrel{(v)}{\leq} 24\sqrt{\frac{2k \log(3k) \tilde{K}}{|S|}} \int_0^1 \sqrt{(2 \text{fat}_{c\gamma}(\mathcal{H}) + 1) \log\left(\frac{2}{\gamma}\right)} d\gamma + 3\sqrt{\frac{\ln(2/\delta)}{2|S|}} \\ &\leq \epsilon \end{aligned}$$

(i) stems from Theorem 5 (generalization error by Rademacher complexity bound for the function class $\text{max-conv}((\mathcal{F}_{\mathcal{H}}^{(j)})_{j \in [k]})$), (ii) stems from Lemma 3, (iii) stems from Theorem 18, (iv) stems from Theorem 12 (fat shattering of max operator) and (v) stems from Lemmas 15 and 16. ■

Proof [of Corollary 10] Let \mathcal{H} be a function class of homogeneous hyperplanes bounded by 1 with domain \mathbb{R}^m .

$$\begin{aligned}
R_n(\max((\mathcal{F}_{\mathcal{H}}^{(j)})_{j \in [k]}))|_{\mathbf{z}} &\leq \inf_{\alpha \geq 0} \left\{ 4\alpha + 12\sqrt{\frac{\tilde{K}}{|S|}} \int_{\alpha}^1 \sqrt{\text{fat}_{c\gamma}(\max((\mathcal{F}_{\mathcal{H}}^{(j)})_{j \in [k]})) \log\left(\frac{2}{\gamma}\right)} d\gamma \right\} \\
&\leq \inf_{\alpha \geq 0} \left\{ 4\alpha + 12\sqrt{\frac{2k \log(3k)\tilde{K}}{|S|}} \int_{\alpha}^1 \sqrt{\max_{j \in [k]} \text{fat}_{c\gamma}(\mathcal{F}_{\mathcal{H}}^{(j)}) \log\left(\frac{2}{\gamma}\right)} d\gamma \right\} \\
&\leq \inf_{\alpha \geq 0} \left\{ 4\alpha + 12\sqrt{\frac{2k \log(3k)\tilde{K}}{|S|}} \int_{\alpha}^1 \sqrt{(2 \text{fat}_{c\gamma}(\mathcal{H}) + 1) \log\left(\frac{2}{\gamma}\right)} d\gamma \right\} \\
&\stackrel{(i)}{\leq} \inf_{\alpha \geq 0} \left\{ 4\alpha + 12c' \sqrt{\frac{2k \log(3k)}{|S|}} \int_{\alpha}^1 \frac{1}{t} \sqrt{\log \frac{2}{t}} dt \right\}
\end{aligned}$$

(i) stems from the bound $\text{fat}_{\delta}(H) \leq \frac{1}{\delta^2}$ (Bartlett and Shawe-Taylor, 1999).
Compute

$$\int_{\alpha}^1 \frac{1}{t} \sqrt{\log \frac{2}{t}} dt = \frac{2}{3} \left(\log(2/\alpha)^{3/2} - (\log 2)^{3/2} \right)$$

and choosing $\alpha = 1/\sqrt{|S|}$ yields

$$\begin{aligned}
R_n(\max((\mathcal{F}_{\mathcal{H}}^{(j)})_{j \in [k]}))|_{\mathbf{z}} &\leq \frac{4}{\sqrt{|S|}} + 8c' \sqrt{\frac{2k \log(3k)}{|S|}} \left(\log(2\sqrt{|S|})^{3/2} - (\log 2)^{3/2} \right) \\
&= O\left(\sqrt{\frac{k \log k \cdot (\log |S|)^3}{|S|}} \right).
\end{aligned}$$

A standard calculation yields sample complexity $m_0 = \mathcal{O}\left(\frac{1}{\epsilon^2} (k \log^2(k/\epsilon) + \log \frac{1}{\delta})\right)$. ■

Acknowledgments

The work of AK and YM was supported in part by grants from the Israel Science Foundation (ISF).

Appendix A.

Proof [of Lemma 8] Suppose that the binary function class $\mathcal{F}_{\mathcal{H}}^{(j)}$ shatters the points $\{(x_1, y_1), \dots, (x_d, y_d)\} \subset \mathcal{X} \times \mathcal{Y}$. That means that for each $b \in \{0, 1\}^d$, there is an $h_b \in \mathcal{H}$ such that $I_{\{h_b(z_j(x_i)) \neq y_i\}} = b_i$ for all $i \in [d]$, where $z_j(x)$ is the j th element in the (ordered) set-valued output of ρ on input x . We claim that \mathcal{H} is able to G -shatter $S = \{z_j(x_1), \dots, z_j(x_d)\} \subset \mathcal{Z}$. Indeed, for each $T \subseteq S$, let $b = b(T) \in \{0, 1\}^S$ be its characteristic function. Taking $f : S \rightarrow \mathcal{Y}$ to be $f(x_i) = y_i$, we see that the definition of G -shattering holds. \blacksquare

Proof [of Lemma 17] To prove the first inequality, observe that for every $m \in \mathbb{N}$,

$$\Pi_{\mathcal{H} \cup \mathcal{G}}(m) \leq \Pi_{\mathcal{H}}(m) + \Pi_{\mathcal{G}}(m),$$

since for every set S , any vector in $\Pi_{\mathcal{H} \cup \mathcal{G}}(S)$ belongs to $\Pi_{\mathcal{H}}(S)$ or $\Pi_{\mathcal{G}}(S)$, where Π denotes the growth function. Taking $m = \text{VC}(\mathcal{H}) + \text{VC}(\mathcal{G}) + 2$ and applying Sauer's Lemma, we get

$$\begin{aligned} \Pi_{\mathcal{H}}(m) + \Pi_{\mathcal{G}}(m) &\leq \sum_{i=0}^{\text{VC}(\mathcal{H})} \binom{m}{i} + \sum_{i=0}^{\text{VC}(\mathcal{G})} \binom{m}{i} \\ &= \sum_{i=0}^{\text{VC}(\mathcal{H})} \binom{m}{i} + \sum_{i=0}^{\text{VC}(\mathcal{G})} \binom{m}{m-i} \\ &= 2^m - \binom{m}{\text{VC}(\mathcal{H}) + 1} \\ &< 2^m, \end{aligned}$$

which shows that $\mathcal{H} \cup \mathcal{G}$ cannot shatter sets of size m and proves the first claim.

To prove the second inequality, observe that

$$\text{fat}_{\gamma}(\mathcal{H} \cup \mathcal{G}) = \text{fat}_{\gamma}^0((\mathcal{H} \cup \mathcal{G}) - r) = \text{fat}_{\gamma}^0((\mathcal{H} - r) \cup (\mathcal{G} - r))$$

for some $r \in \mathbb{R}$. Thus, to prove the claim, it suffices to show that

$$\text{fat}_{\gamma}^0(\mathcal{H} \cup \mathcal{G}) \leq \text{fat}_{\gamma}^0(\mathcal{H}) + \text{fat}_{\gamma}^0(\mathcal{G}) + 1.$$

To this end, we apply the same exact technique as in the proof of Theorem 12. We convert \mathcal{F}, \mathcal{G} and $\mathcal{F} \cup \mathcal{G}$ first to the ternary-valued classes $\mathcal{F}^*, \mathcal{G}^*, (\mathcal{F} \cup \mathcal{G})^* \subseteq \{-\gamma, \gamma, \star\}^X$ and then to binary classes via the ‘‘disambiguation’’ mapping φ furnished by Lemma 14. Clearly,

$$\varphi((\mathcal{F} \cup \mathcal{G})^*) \subseteq \varphi(\mathcal{F}^*) \cup \varphi(\mathcal{G}^*)$$

and $\text{fat}_{\gamma}^0(\mathcal{F}) = \text{VC}(\varphi(\mathcal{F}^*))$, with analogous relations for \mathcal{G} and $\mathcal{F} \cup \mathcal{G}$. It follows that

$$\begin{aligned} \text{fat}_{\gamma}^0(\mathcal{H} \cup \mathcal{G}) &= \text{VC}(\varphi((\mathcal{F} \cup \mathcal{G})^*)) \\ &\leq \text{VC}(\varphi(\mathcal{F}^*) \cup \varphi(\mathcal{G}^*)) \\ &\leq \text{VC}(\varphi(\mathcal{F}^*)) + \text{VC}(\varphi(\mathcal{G}^*)) + 1 \\ &= \text{fat}_{\gamma}^0(\mathcal{F}) + \text{fat}_{\gamma}^0(\mathcal{G}) + 1, \end{aligned}$$

which proves the claim. \blacksquare

References

- Dana Angluin and Philip Laird. Learning from noisy examples. *Machine Learning*, 2(4):343–370, Apr 1988. ISSN 1573-0565. doi: 10.1023/A:1022873112823. URL <https://doi.org/10.1023/A:1022873112823>.
- Peter Bartlett and John Shawe-Taylor. *Generalization performance of support vector machines and other pattern classifiers*, pages 43–54. MIT Press, Cambridge, MA, USA, 1999. ISBN 0-262-19416-3.
- Peter L. Bartlett and Shahar Mendelson. Rademacher and gaussian complexities: Risk bounds and structural results. *Journal of Machine Learning Research*, 3:463–482, 2002.
- Shai Ben-David, Nicolò Cesa-Bianchi, David Haussler, and Philip M. Long. Characterizations of learnability for classes of $\{0, \dots, n\}$ -valued functions. *J. Comput. Syst. Sci.*, 50(1):74–86, 1995.
- A. Ben-Tal, L. El Ghaoui, and A.S. Nemirovski. *Robust Optimization*. Princeton Series in Applied Mathematics. Princeton University Press, October 2009.
- Anselm Blumer, Andrzej Ehrenfeucht, David Haussler, and Manfred K. Warmuth. Learnability and the Vapnik-Chervonenkis dimension. *J. Assoc. Comput. Mach.*, 36(4):929–965, 1989. ISSN 0004-5411.
- Stéphane Boucheron, Olivier Bousquet, and Gábor Lugosi. Theory of classification: A survey of recent advances. *ESAIM Probab. Statist.*, 9:323–375, 2005.
- Nicolò Cesa-Bianchi, Yishay Mansour, and Gilles Stoltz. Improved second-order bounds for prediction with expert advice. In *COLT*, volume 3559 of *Lecture Notes in Computer Science*, pages 217–232. Springer, 2005.
- Monika Csikos, Andrey Kupavskii, and Nabil H. Mustafa. Optimal bounds on the vc-dimension. *CoRR*, abs/1807.07924, 2018.
- Ofer Dekel and Ohad Shamir. Learning to classify with missing and corrupted features. In *Proceedings of the 25th International Conference on Machine Learning, ICML '08*, pages 216–223, New York, NY, USA, 2008. ACM. ISBN 978-1-60558-205-4. doi: 10.1145/1390156.1390184. URL <http://doi.acm.org/10.1145/1390156.1390184>.
- R.M Dudley. The sizes of compact subsets of hilbert space and continuity of gaussian processes. *Journal of Functional Analysis*, 1(3):290 – 330, 1967. ISSN 0022-1236. doi: [http://dx.doi.org/10.1016/0022-1236\(67\)90017-1](http://dx.doi.org/10.1016/0022-1236(67)90017-1). URL <http://www.sciencedirect.com/science/article/pii/0022123667900171>.
- Uriel Feige, Yishay Mansour, and Robert E. Schapire. Learning and inference in the presence of corrupted inputs. In *Proceedings of The 28th Conference on Learning Theory, COLT 2015, Paris, France, July 3-6, 2015*, pages 637–657, 2015.
- Uriel Feige, Yishay Mansour, and Robert E. Schapire. Robust inference for multiclass classification. In *Algorithmic Learning Theory, ALT 2018, 7-9 April 2018, Lanzarote, Canary Islands, Spain*, pages 368–386, 2018.

- Yoav Freund and Robert E. Schapire. Adaptive game playing using multiplicative weights. *Games and Economic Behavior*, 29(1-2):79–103, 1999. URL <https://EconPapers.repec.org/RePEc:eee:gamebe:v:29:y:1999:i:1-2:p:79-103>.
- Amir Globerson and Sam Roweis. Nightmare at test time: Robust learning by feature deletion. In *Proceedings of the 23rd International Conference on Machine Learning, ICML '06*, pages 353–360, New York, NY, USA, 2006. ACM. ISBN 1-59593-383-2. doi: 10.1145/1143844.1143889. URL <http://doi.acm.org/10.1145/1143844.1143889>.
- Ian Goodfellow, Jonathon Shlens, and Christian Szegedy. Explaining and harnessing adversarial examples. In *International Conference on Learning Representations*, 2015. URL <http://arxiv.org/abs/1412.6572>.
- Lee-Ad Gottlieb, Aryeh Kontorovich, and Robert Krauthgamer. Efficient classification for metric data. *IEEE Transactions on Information Theory*, 60(9):5750–5759, 2014. doi: 10.1109/TIT.2014.2339840. URL <http://dx.doi.org/10.1109/TIT.2014.2339840>.
- Michael Kearns and Ming Li. Learning in the presence of malicious errors. *SIAM Journal on Computing*, 22:807–837, 1993.
- Aryeh Kontorovich. Rademacher complexity of k-fold maxima of hyperplanes. 2018.
- Alexey Kurakin, Ian J. Goodfellow, and Samy Bengio. Adversarial machine learning at scale. 2017. URL <https://arxiv.org/abs/1611.01236>.
- Pavel Laskov and Richard Lippmann. Machine learning in adversarial environments. *Mach. Learn.*, 81(2):115–119, November 2010. ISSN 0885-6125. doi: 10.1007/s10994-010-5207-6. URL <https://doi.org/10.1007/s10994-010-5207-6>.
- Aleksander Madry, Aleksandar Makelov, Ludwig Schmidt, Dimitris Tsipras, and Adrian Vladu. Towards deep learning models resistant to adversarial attacks. *CoRR*, abs/1706.06083, 2017.
- Yishay Mansour, Aviad Rubinfeld, and Moshe Tennenholtz. Robust probabilistic inference. In *Proceedings of the Twenty-Sixth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2015, San Diego, CA, USA, January 4-6, 2015*, pages 449–460, 2015.
- S. Mendelson and R. Vershynin. Entropy and the combinatorial dimension. *Invent. Math.*, 152(1):37–55, 2003. ISSN 0020-9910. doi: 10.1007/s00222-002-0266-3. URL <http://dx.doi.org/10.1007/s00222-002-0266-3>.
- Mehryar Mohri, Afshin Rostamizadeh, and Ameet Talwalkar. *Foundations Of Machine Learning*. The MIT Press, 2012.
- S. Moosavi-Dezfooli, A. Fawzi, and P. Frossard. Deepfool: A simple and accurate method to fool deep neural networks. In *2016 IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, pages 2574–2582, June 2016. doi: 10.1109/CVPR.2016.282.
- Ludwig Schmidt, Shibani Santurkar, Dimitris Tsipras, Kunal Talwar, and Aleksander Madry. Adversarially robust generalization requires more data. *CoRR*, abs/1804.11285, 2018.

- Choon H. Teo, Amir Globerson, Sam T. Roweis, and Alex J. Smola. Convex learning with invariances. In J. C. Platt, D. Koller, Y. Singer, and S. T. Roweis, editors, *Advances in Neural Information Processing Systems 20*, pages 1489–1496. Curran Associates, Inc., 2008. URL <http://papers.nips.cc/paper/3218-convex-learning-with-invariances.pdf>.
- Florian Tramèr, Nicolas Papernot, Ian J. Goodfellow, Dan Boneh, and Patrick D. McDaniel. The space of transferable adversarial examples. *CoRR*, abs/1704.03453, 2017.
- L. G. Valiant. Learning disjunction of conjunctions. In *Proceedings of the 9th International Joint Conference on Artificial Intelligence - Volume 1*, IJCAI'85, pages 560–566, San Francisco, CA, USA, 1985. Morgan Kaufmann Publishers Inc. ISBN 0-934613-02-8. URL <http://dl.acm.org/citation.cfm?id=1625135.1625242>.